

Holomorphic curves in exploded manifolds: compactness

Brett Parker
bparker@msri.org

November 11, 2009

Abstract

This paper establishes compactness results for the moduli stack of holomorphic curves in suitable exploded manifolds. This result together with the analysis in [8] allows the definition of Gromov Witten invariants of these exploded manifolds.

Contents

1	Introduction	1
2	Strict tamings	4
3	Estimates	6
4	Compactness	27
A	Equivariant coordinates	40
B	A symplectic analogue of the explosion functor	43
C	Construction of a strict taming	45

1 Introduction

This paper establishes compactness results for the moduli stack of holomorphic curves in exploded manifolds. An introduction to exploded manifolds may be found in [7].

The topology in which we shall establish compactness is the $C^{\infty,1}$ topology, which is both a topology and a level of regularity. A $C^{\infty,1}$ function can be viewed as a generalization of a function on a manifold with cylindrical ends which is smooth and has exponential convergence at cylindrical ends. In [8] it is shown that the moduli stack of holomorphic curves has regularity $C^{\infty,1}$ in the sense that in a neighborhood of any holomorphic curve for which the $\bar{\partial}$ equation is transversal, the moduli stack of holomorphic curves can be represented by $C^{\infty,1}$ family of holomorphic curves. More generally, a virtual moduli space is constructed in [8] which is locally constructed using $C^{\infty,1}$ families of curves.

The compactness established in this paper allows us to know that components of this virtual moduli space are compact, and define Gromov Witten invariants for suitable exploded manifolds.

The topology in which the moduli space of curves is compact seems unnatural from the perspective of differential topology because of bubbling phenomena and node formation. Similar phenomena occur in the category of exploded manifolds, but unlike the smooth category, all the bubbling and node formation behavior can happen within smooth (or $C^{\infty,1}$) connected families of maps. Still, in talking about a converging sequence of holomorphic curves, we must deal with the fact that the domain of the curves may change.

We shall say that a sequence of $C^{\infty,1}$ curves

$$f_i : (\mathbf{C}, j_i) \longrightarrow \mathbf{B}$$

converges to a $C^{\infty,1}$ curve f if the following holds:

There exists a sequence of families of $C^{\infty,1}$ curves,

$$\begin{array}{ccc} (\hat{\mathbf{C}}, j_i) & \xrightarrow{\hat{f}^i} & \hat{\mathbf{B}} \\ \downarrow & & \\ \mathbf{F} & & \end{array}$$

and a sequence of points $p_i \in \mathbf{F}$ so that

1. The fiber of $(\hat{\mathbf{C}}, j_i)$ over p^i is $(\mathbf{C}, j_i)$, and the restriction of $\hat{f}^i$ to the fiber over p^i is f^i .
2. This sequence of families converges in $C^{\infty,1}$ to the family

$$\begin{array}{ccc} (\hat{\mathbf{C}}, j) & \xrightarrow{\hat{f}} & \hat{\mathbf{B}} \\ \downarrow & & \\ \mathbf{F} & & \end{array}$$

in the sense that the fiberwise complex structures j_i converge in $C^{\infty,1}$ to j and the maps $\hat{f}^i$ converge in $C^{\infty,1}$ to $\hat{f}$

3. The sequence of points p^i in $\mathbf{F}$ converge to some point $p \in F$ so that f is given by the restriction of $\hat{f}$ to the fiber over p .

One scary aspect of this definition is that the corresponding topology on the moduli stack of holomorphic curves is non-Hausdorff in the same way that the topology on an exploded manifold is non-Hausdorff. In particular all the points $p \in F$ for which $p_i \rightarrow p$ correspond to limits of our sequence of curves. This is to be expected as we want to view the moduli stack of holomorphic curves as analogous to an exploded manifold rather than simply as a topological space.

The almost complex exploded manifolds $(\mathbf{B}, J)$ for which we prove compactness for the moduli space of holomorphic curves satisfy the following assumptions:

1. $\mathbf{B}$ is basic. This assumption roughly says that the tropical part of $\mathbf{B}$ is a union of convex polytopes glued along faces so that any two polytopes are glued in at most one way.

2. $\mathbf{B}$ is complete. This plays the role that compactness plays in the case of smooth manifolds. This means that $\mathbf{B}$ is compact as a topological space, and that the tropical part of $\mathbf{B}$ is a union of complete polytopes.
3. The almost complex structure J on $\mathbf{B}$ is civilized. This means that J induces an almost complex structure on the smooth part of $\mathbf{B}$, which is helpful because it allows us to use standard holomorphic curve results such as the removable singularity theorem in our setting. There is no obstruction to modifying an almost complex structure to make it civilized.
4. There is a strict taming Ω of J . Strict tamings are discussed in section 2. A strict taming plays the role that a symplectic structure plays in the case of smooth manifolds. Each strict taming is a set Ω of closed two forms ω on $\mathbf{B}$ which are non negative on holomorphic planes. Given any holomorphic curve $f : \mathbf{C} \longrightarrow \mathbf{B}$, the integral of $f^*\omega$ does not depend on the choice of $\omega \in \Omega$. This result of this integral is called the Ω energy of the holomorphic curve f .

With these assumptions and the $C^{\infty,1}$ topology understood, we can state a compactness theorem as follows:

Given a basic, complete exploded manifold $\mathbf{B}$ with a civilized almost complex structure J and strict taming Ω , the moduli stack $\mathcal{M}_{g,n,E}(\mathbf{B})$ of stable holomorphic curves with a fixed genus g and number of punctures n , and with Ω energy less than E is compact using the $C^{\infty,1}$ topology.

A more general version involving the case of curves in a family of exploded manifolds is stated on page 27. This compactness result can be viewed as a generalization of the compactness results for holomorphic found in [2], [4], [5], and [3].

The following are some examples of almost complex exploded manifolds satisfying the requirements of the above theorem.

- Any compact manifold with an almost complex structure tamed by a symplectic form will satisfy the above assumptions.
- The explosion $\text{Expl } M$ of any complex manifold M with normal crossing divisors so that each divisor is an embedded submanifold will be basic. If M is compact and there is a symplectic form on M which tames the complex structure, then $\text{Expl } M$ will satisfy the above assumptions. This case is useful for Gromov Witten invariants relative to normal crossing divisors.
- Given any compact symplectic manifold M with orthogonally intersecting codimension 2 symplectic submanifolds, we can construct an exploded manifold $\mathbf{M}$ using a similar construction to the explosion functor. There exists an almost complex structure J on $\mathbf{M}$ satisfying the above assumptions. This case is useful to define Gromov Witten invariants relative to these symplectic submanifolds.

2 Strict tamings

In this section, we define a strict taming of a civilized almost complex structure on an exploded manifold. A strict taming will play the role that a symplectic form usually plays.

Consider the special case of a basic, complete, exploded manifold $\mathbf{B}$ so that each polytope in the tropical part of $\mathbf{B}$ is either a standard simplex or quadrant $[0, \infty)^n$. In this case the smooth part of $\mathbf{B}$ is a stratified space, where the closure of each strata $[\bar{\mathbf{B}}_i]$ is a compact manifold. If there is a civilized almost complex structure J on $\mathbf{B}$, then $[\bar{\mathbf{B}}_i]$ is an almost complex manifold, where lower dimensional strata appear as almost complex submanifolds. In this case, a natural candidate for a taming form is a symplectic form ω which tames J on each $[\bar{\mathbf{B}}_i]$. Lemma C.1 shows that if J is integrable, a strict taming of J can be constructed from ω . If J is not integrable, it may be modified so that a strict taming of the modified J can be constructed from ω .

More generally, the smooth part of $\mathbf{B}$ will be a stratified space in which the closure of strata $[\bar{\mathbf{B}}_i]$ are singular almost complex manifolds. We shall now give a definition of a symplectic form on $\mathbf{B}$ which tames a civilized almost complex structure J , which roughly speaking gives a symplectic form taming J on each $[\bar{\mathbf{B}}_i]$.

Definition 2.1. *Given an exploded manifold $\mathbf{B}$ with a civilized almost complex structure J , say that a two form ω on $\mathbf{B}$ is a symplectic form which tames J if each point in $\mathbf{B}$ has an open neighborhood U with a smooth map*

$$\iota : U \rightarrow \mathbb{R}^{2n}$$

so that

1. every smooth function on U is the pullback of some smooth function on $\mathbb{R}^n$
2. there exists some almost complex structure $\hat{J}$ on $\mathbb{R}^{2n}$ so that $\hat{J}d\iota = d\iota J$
3. there exists some symplectic form $\hat{\omega}$ on $\mathbb{R}^{2n}$ which is positive on $\hat{J}$ -holomorphic planes so that $\omega = \iota^*\hat{\omega}$

The point of using civilized almost complex structures and the above type of taming forms is that it becomes easy to apply known results such as the removable singularity theorem and the monotonicity lemma from the analysis of holomorphic curves in smooth manifolds.

Example 2.2.

Recall from [7] that the explosion functor constructs an exploded manifold $\text{Expl } M$ from a complex manifold with normal crossing divisors M . If ω is any symplectic form on M which tames the complex structure, then the pullback of ω to $\text{Expl } M$ via the smooth part map $\text{Expl } M \rightarrow M$ is a symplectic form on $\text{Expl } M$ that tames the complex structure.

The definition of a symplectic taming form is clearly inadequate for taming holomorphic curves in $\mathbf{T}^n$ where it is satisfied trivially by a two form that is

identically 0 because all smooth functions on $\mathbf{T}^n$ are constant. We remedy this by defining a strict taming. (Note that we shall use freely the fact from [7] that any holomorphic curve in $\mathbf{B}$ lifts to a holomorphic curve (with a refined domain) in $\mathbf{B}'$ for any refinement $\mathbf{B}' \longrightarrow \mathbf{B}$.)

Definition 2.3. *A strict taming of a civilized almost complex structure J on a complete exploded manifold $\mathbf{B}$ is a set Ω of closed two forms on refinements of $\mathbf{B}$ containing at least one symplectic form which tames J . Each form in Ω must be nonnegative on J holomorphic planes, and there must exist a metric g on $\mathbf{B}$ and a positive number $c > 0$ smaller than the injectivity radius of g so that*

1. *Given any complete smooth curve $f : \mathbf{C} \longrightarrow \mathbf{B}$, the ω energy, $\int_{\mathbf{C}} f^* \omega$ is a constant independent of $\omega \in \Omega$*
2. *Given any point $p \in \mathbf{B}$, there exists a taming form $\omega_p \in \Omega$ so that on the ball of radius c around p ,*

$$\omega_p(v, Jv) \geq g(v, v)$$

The Ω energy of a map $f : \mathbf{C} \longrightarrow \mathbf{B}$ where $\mathbf{C}$ is two dimensional and oriented is given by

$$E_{\Omega}(f) := \sup_{\omega \in \Omega} \int f^* \omega$$

For example, any compact almost complex manifold tamed by a symplectic form is strictly tamed by that symplectic form. A strict taming is what is needed to get local area bounds on holomorphic curves. Strict tamings can be viewed as a generalization of the notion of energy used to tame holomorphic curves in symplectic field theory [1].

Example 2.4.

A simple example of a strict taming is given by the case of the explosion of a compact toric manifold M relative to its toric boundary divisors. We can regard $\text{Expl } M$ as a refinement of $\mathbf{T}^n$ given by subdividing the tropical part of $\mathbf{T}^n$ using the toric fan of M . There are global holomorphic coordinates $(\tilde{z}_1, \dots, \tilde{z}_n)$ on $\text{Expl } M$. Choose some symplectic form ω on M which tames the complex structure. Using the standard flat metric in which the real and imaginary parts of $\tilde{z}_i \frac{\partial}{\partial \tilde{z}_i}$ form an orthogonal basis, on a ball of radius 1 around $(1t^0, \dots, 1t^0)$, ω is bounded below on holomorphic planes. Therefore, if g indicates the standard flat metric scaled by some small positive constant,

$$w(v, Jv) \geq g(v, v)$$

on a ball of radius 1 around $(1t^0, \dots, 1t^0)$.

Given any other point $p = (c_1 t^{a_1}, \dots, c_n t^{a_n})$, consider the refinement $\mathbf{M}'$ of $\text{Expl } M$ given by subdividing $\text{Expl } M$ by the toric fan of M translated to be centered at p . We may use the same global coordinates $(\tilde{z}_1, \dots, \tilde{z}_n)$ on $\mathbf{M}'$. Now there is a smooth map

$$\phi_p : \mathbf{M}' \longrightarrow \text{Expl } M$$

given by

$$(\tilde{z}_1, \dots, \tilde{z}_n) \mapsto (c_1^{-1} t^{-a_1} \tilde{z}_1, \dots, c_n t^{-a_n} \tilde{z}_n)$$

We can now define $\omega_p := \phi_p^* \omega$. The map ϕ_p preserves J and the metric g , so ω_p is non negative on holomorphic planes and satisfies

$$w_p(v, Jv) \geq g(v, v)$$

on a ball of radius 1 around p . The integral of ω_p on any complete exploded curve will be the same as the integral of ω , so the set $\Omega := \{\omega_p \text{ for all } p \in \text{Expl } M\}$ is a strict taming.

Using the construction of Lemma C.1 we can construct a strict taming Ω in the following cases:

- If $\text{Expl } M$ is the explosion of a compact complex manifold with normal crossing divisors which are embedded submanifolds, and ω is a symplectic form on M taming the complex structure, there is a strict taming Ω of the complex structure on $\text{Expl } M$ which contains ω .
- Example B.1 on page 43 constructs an exploded manifold $\mathbf{M}$ from a compact symplectic manifold M with orthogonally intersecting codimension 2 symplectic submanifolds. This construction is similar to the construction of the explosion functor, so the smooth part of $\mathbf{M}$ is M , and the tropical part of $\mathbf{M}$ is the dual intersection complex which has a vertex for each connected component of M , a ray for each of our codimension 2 submanifolds, and an n -dimensional face $[0, \infty)^n$ for each n -fold intersection. There exists an almost complex structure on $\mathbf{M}$ with a strict taming Ω which contains the pullback of the symplectic form from M .
- Let $\mathbf{B}$ is a basic exploded manifold so that each polytope in the tropical part of $\mathbf{B}$ is a standard simplex. Suppose that J is an almost complex structure on $\mathbf{B}$ so that for each exploded function $\tilde{z}$ locally defined on $\mathbf{B}$, $\tilde{z}^{-1} \bar{\partial} \tilde{z}$ is a smooth function. (For example any integrable complex structure satisfies this condition.) If ω is a symplectic form on $\mathbf{B}$ which tames J , then there exists a strict taming of J which contains ω .

If J is a civilized almost complex structure on $\mathbf{B}$ tamed by ω which does not satisfy the above condition, then it may be modified to an almost complex structure still tamed by ω which does, and we may then construct a strict taming which contains the new almost complex structure.

3 Estimates

In this section, we shall prove the analytic estimates required for our compactness theorem. These estimates do not differ much from the standard estimates used to study (pseudo)holomorphic curves in smooth symplectic manifolds. They are also proved using roughly the same methods as in the smooth case. We shall make the following assumptions:

1. We'll assume that our target $\mathbf{B}$ is complete. This is required to give us the kind of bounded geometry found in the smooth case when the target is compact or has a cylindrical end. We shall also assume that $\mathbf{B}$ is basic. The assumption that $\mathbf{B}$ is basic is mainly for convenience in the arguments that follow. It is most useful in establishing coordinates and notation for the construction of families of smooth curves in the proof of Theorem 4.1.

2. We'll assume our almost complex structure J on $\mathbf{B}$ is civilized. This artificial (but easy to achieve) assumption means that J induces a smooth almost complex structure on the smooth parts of $\mathbf{B}$. This allows us to use standard estimates from the study of holomorphic curves in smooth manifolds with minimal modification. This assumption is needed to have holomorphic curves be smooth maps instead of just $C^{\infty,1}$ maps.
3. We'll assume that J has a strict taming Ω . This is required to get a local area bound for holomorphic curves. We shall also work with a particular symplectic form $\omega \in \Omega$. This together with J defines a pseudo metric on $\mathbf{B}$ which collapses all tropical directions (so it is like a metric on the smooth part $[\mathbf{B}]$.) The strategy of proof for most of our estimates is roughly as follows: first prove a weak estimate in ω 's pseudo metric, and then improve this to get an estimate in an actual metric on $\mathbf{B}$. We shall use this strategy to prove estimates for the the derivative at the center of holomorphic disks with bounded Ω energy and small ω energy, and to get strong estimates on the behavior of holomorphic cylinders with bounded Ω energy and small ω energy.

This section ends with Proposition 3.18, which is a careful statement of the fact that any holomorphic curve with bounded Ω energy, genus, and number of punctures can be decomposed into a bounded number of components, which either have 'bounded conformal structure and bounded derivative', or are annuli with small ω energy, (and so we have strong estimates on their behavior.)

Throughout this section, we shall use the notation $(\mathbf{B}, J, \Omega, \omega, g)$ to indicate a smooth, basic, complete exploded manifold $\mathbf{B}$, a civilized almost complex structure J , a strict taming Ω , a choice of symplectic taming form $\omega \in \Omega$, and a metric g . If we say that a constant depends continuously on $(\mathbf{B}, J, \Omega, \omega, g)$, we mean that if we have a family $\hat{\mathbf{B}} \rightarrow \mathbf{G}$ with such a structure on each fiber, the constant can be chosen a continuous function on $\mathbf{G}$.

We shall use the following lemma which bounds the geometry of a complete exploded manifold. Note that given a metric g on some complete $\mathbf{B}$, the injectivity radius is a continuous function on $\mathbf{B}$, and is therefore bounded below.

Lemma 3.1. *Given a complete exploded manifold $\mathbf{B}$ with a metric g , there exists some radius $R > 0$ smaller than the injectivity radius of $(\mathbf{B}, g)$ so that the following holds:*

Given a sequence of points $p_n \in \mathbf{B}$ converging to p , consider the ball of radius R around p_n as a smooth manifold $B_R(p_n)$.

There exists a sequence of diffeomorphisms

$$f_n : B_R(p_n) \rightarrow B_R(p)$$

so that

$$(f_n)_*g \text{ converges to } g \text{ in } C^\infty,$$

and given any smooth tensor field θ on $\mathbf{B}$ (such as an almost complex structure), $(f_n)_(\theta)$ converges to θ in C^∞ .*

Similarly, given a family $\hat{\mathbf{B}} \rightarrow \mathbf{G}$ of complete exploded manifolds with fiberwise metrics g , we may choose $R > 0$ to depend continuously on $\mathbf{G}$ so that the following holds:

*Given a sequence of points $p_n \in \hat{\mathbf{B}}$ converging to p , consider the ball of radius R around p_n in the fiber containing p_n as a smooth manifold $B_R(p_n)$. Then the above conclusion holds: There is a sequence of diffeomorphisms $f_n : B_R(p_n) \rightarrow B_R(p)$ so that $(f_n)_*g$ converges to g in C^∞ , and given any smooth vertical tensor field θ on $\hat{\mathbf{B}}$, $(f_n)_*(\theta)$ converges to θ in C^∞ .*

Proof: By choosing R small enough, we may assume that $B_R(p)$ is contained in a single coordinate chart, and $B_R(p_n)$ is eventually contained in the same coordinate chart. The lemma then follows from the fact that all smooth tensor fields (and their derivatives) can be locally written as some sum of smooth functions on $\mathbf{B}$ times some basis tensor fields. $\square$

This tells us that the geometry of a complete exploded manifold is bounded in the same way that the geometry of a compact manifold is bounded. The analogous result for families is also true.

The following is a consequence of our bounded geometry and the standard monotonicity lemma for holomorphic curves first proved in [2]. (The proof involves a bound on the covariant derivatives of J , the curvature of g and the injectivity radius, all of which vary continuously in families.)

Lemma 3.2. *Given $(\mathbf{B}, J, \Omega, g)$, for any $\epsilon > 0$*

there exists some $E > 0$ depending continuously on $(\mathbf{B}, J, \Omega, g)$

so that any non constant J holomorphic curve

which passes through a point p

and which is a complete map when restricted to the ϵ ball around p

has Ω energy greater than E .

Proof: Let R be as in Lemma 3.1. Consider a holomorphic curve passing through p which is proper when restricted to the ball of radius $\min\{R, \epsilon\}$ around p . The monotonicity lemma for holomorphic curves proved in [2] implies that there is some lower bound $A > 0$ for the area of our holomorphic curve inside our ball. This area lower bound depends continuously on R , ϵ , the covariant derivatives of J and the curvature of g . The bounded geometry from Lemma 3.1 therefore implies that there is a universal area lower bound $A > 0$ depending continuously on $(\mathbf{B}, J, g)$ so that any nonconstant holomorphic curve which passes through any point p and which is proper when restricted to the ϵ ball around p has area bounded below by A inside $B_\epsilon(p)$. Given a strict taming Ω , we may convert this lower bound for area in to a lower bound for the Ω energy of the holomorphic curve. $\square$

Definition 3.3. Given $(\mathbf{B}, J, \omega)$, define the pseudo-metric

$$\langle v, w \rangle_\omega := \frac{1}{2}(\omega(v, Jw) + \omega(w, Jv))$$

This defines a distance function d_ω on $[\mathbf{B}]$.

Lemma 3.4. Given $(\mathbf{B}, J, \Omega, \omega)$ and a choice of $\epsilon > 0$,

there exists some E depending continuously on $(\mathbf{B}, J, \Omega, \omega)$

so that any non constant J holomorphic curve

which passes through a point p

and which is a complete map when restricted to the d_ω -ball of size ϵ around p

has ω energy greater than E .

Proof:

The fact that $\mathbf{B}$ is complete means in particular that it can be covered by a finite number of coordinate charts, $\mathbb{R}^{2n} \times \mathbf{T}_P^m$. These coordinate charts $\mathbb{R}^{2n} \times \mathbf{T}_P^m$ with J and ω can be regarded as subsets of more standard charts $\mathbb{R}^{2n} \times (\mathbf{T}_1^1)^a \times \mathbf{T}^b$ cut out by setting some monomials equal to 1. This has the advantage that the smooth part of this is just equal to $\mathbb{R}^{2n} \times \mathbb{C}^a$, and we can assume our symplectic and almost complex structures just come from smooth ones here (this uses that J is civilized), so we can use standard holomorphic curve results. Note also that the above pseudo metric is actually a metric on this smooth part. The size of covariant derivatives of J in this metric give continuous functions on $\mathbf{B}$, and are therefore bounded because $\mathbf{B}$ is complete. For a given point $q \in [U]$ in the smooth part of a coordinate chart U , the usual monotonicity lemma from [2] implies that there exists some lower bound $E_U(q) > 0$ for the ω energy of holomorphic curves which are non constant in $[U]$, pass through $q \in [U]$ and complete when restricted to the d_ω -ball of size ϵ around $q \in [U]$. This lower bound $E_U(q)$ can be chosen to depend continuously on $q \in [U]$. Cover $\mathbf{B}$ by a finite number of coordinate charts U_i . The function $E(q) := \max_i E_{U_i}(q)$ is positive and lower semicontinuous on $[\mathbf{B}]$, which is compact. Therefore, there exists some lower bound $E > 0$ for $E(q)$. In the case of a family $\hat{\mathbf{B}} \rightarrow \mathbf{G}$, cover $\hat{\mathbf{B}}$ by coordinate charts so that each fiber is only contained in a finite number of coordinate charts, then $E(q)$ has a lower bound E which can be chosen to depend continuously on $\mathbf{G}$.

Suppose now that we have a holomorphic curve that passes through a point p and which is complete when restricted to the d_ω -ball of size ϵ around p . Then either our holomorphic curve must have energy greater than E or it must have some connected component which is constant in $[B]$. This connected component must therefore not touch the boundary of our ϵ ball, and must be a complete map to $\mathbf{B}$. The fact that Ω is a strict taming then allows us to use Lemma 3.2 to complete our proof. $\square$

The above proof also implies the following:

Lemma 3.5. Given $(\mathbf{B}, J, \Omega, \omega)$,

there exists some $E > 0$ which can be chosen to depend continuously on $(\mathbf{B}, J, \Omega, \omega)$

so that given any complete holomorphic curve in $\mathbf{B}$,

the ω energy of any smooth component is either greater than E , or the image of that smooth component in $[B]$ is a point.

Lemma 3.6. *Given $(\mathbf{B}, J, \Omega, \omega)$, for any $\epsilon > 0$,*

there exists some $E > 0$ depending continuously on $(\mathbf{B}, J, \Omega, \omega)$

so that any J holomorphic map f of $\{e^{-(R+1)} < |z| < e^{(R+1)}\} \subset \mathbb{C}$ with ω -energy less than E

is contained inside a d_ω -ball of radius ϵ on the smaller annulus

$$\{e^{-R} \leq |z| \leq e^R\}$$

Proof:

Suppose to the contrary that this lemma is false. All distances in the following shall use d_ω , ω 's pseudo metric.

Choose E small enough so that Lemma 3.4 holds for $\frac{\epsilon}{8}$ balls. There must be a path in $\{e^{-(R+1)} < |z| < e^{(R+1)}\}$ joining $z = 1$ with an end of this annulus contained entirely inside the ball $B_{\frac{\epsilon}{8}}(f(1))$. (Otherwise, f restricted to some subset would be a proper map to $B_{\frac{\epsilon}{8}}(f(1))$, and we could use Lemma 3.4 to get a contradiction.) Suppose without loss of generality that it connects 1 with the circle $|z| = e^{-(R+1)}$.

Suppose for a second that there exists some point z so that $e^{-(R+\frac{3}{4})} < |z| < e^{-(R+\frac{1}{4})}$ and $f(z) \notin B_{\frac{3\epsilon}{8}}(f(1))$. As above, there must exist some path connecting z with an end which is contained in the ball $B_{\frac{\epsilon}{8}}(f(z))$. There must therefore be some region conformal to $[0, \pi] \times [0, \frac{1}{4}]$ so that the $d_\omega(f(0, t), f(\pi, t)) > \frac{\epsilon}{8}$. Then the Cauchy-Schwartz inequality tells us that

$$\int_0^\pi \int_0^{\frac{1}{4}} |df|_\omega^2 > \frac{1}{4} \frac{(\frac{\epsilon}{8})^2}{\pi}$$

This in turn gives a uniform lower bound for the energy of our curve. This case can therefore be discarded, and we can assume that all z satisfying $e^{-(R+\frac{3}{4})} < |z| < e^{-(R+\frac{1}{4})}$ are contained in $B_{\frac{3\epsilon}{8}}(f(1))$.

As we are assuming this lemma is false, there must be some point z_2 so that $e^{-R} \leq |z_2| \leq e^R$ and $f(z_2) \notin B_\epsilon(f(1))$. Repeating the above argument (with z_2 in place of the point $z = 1$) gives us that z must be contained in $B_{\frac{3\epsilon}{8}}(f(z_2))$ when $e^{R+\frac{1}{4}} \leq |z| \leq e^{R+\frac{3}{4}}$.

We then have f contained in balls which are at least $\frac{\epsilon}{4}$ apart on the boundary of an annulus, thus we can apply Lemma 3.4 to some point in the image under f of the interior at least $\frac{\epsilon}{8}$ from both balls to obtain a lower bound for the energy and a contradiction to the assumption that the lemma was false. $\square$

The following gives useful coordinates for the analysis of holomorphic curves:

Lemma 3.7. *Given $(\mathbf{B}, J, \Omega, \omega)$,*

there exist constants c_i and $\epsilon > 0$ depending continuously on $(\mathbf{B}, J, \Omega, \omega)$

so that for all points $p \in \mathbf{B}$,

there exists some coordinate neighborhood U of p

so that the smooth part, $[U]$ can be identified with a relatively closed subset of the open ϵ ball in $\mathbb{C}^n$ with almost complex structure $\hat{J}$ and some flat $\hat{J}$ preserving connection ∇

so that

1. *p is sent to 0, and the restriction of $\hat{J}$ to $[U]$ is J .*
2. *The metric $\langle \cdot, \cdot \rangle_\omega$ on $[U]$ is close to the standard flat metric on $\mathbb{C}^n$ so that*

$$\frac{1}{2} \langle v, v \rangle_\omega \leq |v|^2 \leq 2 \langle v, v \rangle_\omega$$

3. *$\hat{J}$ at the point $z \in \mathbb{C}^n$ converges to the standard complex structure J_0 on $\mathbb{C}^n$ as $z \rightarrow 0$ in the sense that*

$$\|\hat{J} - J_0\| \leq c_0 |z|$$

4. *The torsion tensor*

$$\mathbf{T}_\nabla(v, w) := \nabla_v w - \nabla_w v - [v, w]$$

is bounded by c_0 , and has its k th derivatives bounded by c_k .

The only point in this lemma which does not follow from the definition of a civilized almost complex structure and calculation in local coordinates is the fact that the constants involved do not have to depend on the point p . This follows from the fact that $\mathbf{B}$ is complete.

Lemma 3.8. *Given a holomorphic map f of the unit disk to a space with a flat J preserving connection ∇ , the partial derivative of f in the x direction f_x defines a map to $\mathbb{C}^n$ defined by parallel transporting f_x back to the tangent space at $f(0)$ which we then identify with $\mathbb{C}^n$. Such an f_x satisfies the following equation involving the torsion tensor of ∇ , $\mathbf{T}_\nabla$.*

$$\bar{\partial} f_x = \frac{1}{2} J \mathbf{T}_\nabla(f_y, f_x)$$

Proof:

$$\begin{aligned} \bar{\partial} f_x &= \frac{1}{2} (\nabla_{f_x} f_x + J \nabla_{f_y} f_x) \\ &= \frac{1}{2} \nabla_{f_x} (f_x + J f_y) + \frac{1}{2} J (\nabla_{f_y} f_x - \nabla_{f_x} f_y) \\ &= \frac{1}{2} J \mathbf{T}_\nabla(f_y, f_x) \end{aligned}$$

□

As the $\bar{\partial}$ above is the standard linear $\bar{\partial}$ operator, this expression is good for applying the following standard elliptic regularity lemma for the linear $\bar{\partial}$ equation. (An exposition of the proof can be found in [6]). This allows us to get bounds on higher derivatives from bounds on the first derivative of holomorphic functions.

Lemma 3.9. *For a given number k and $1 < p < \infty$, there exists a constant c so that given any map f from the unit disk $D(1)$,*

$$\|f\|_{L_{k+1}^p(D(\frac{1}{2}))} \leq c \left(\|\bar{\partial}f\|_{L_k^p(D(1))} + \|f\|_{L_k^p(D(1))} \right)$$

Lemma 3.10. *Given $(\mathbf{B}, J, \omega)$,*

there exists

some energy E ,

some distance $r > 0$,

and a constant c ,

each depending continuously on $(\mathbf{B}, J, \omega)$

so that any holomorphic map f of a disk $\{|z| \leq 1\}$ into $\mathbf{B}$ which either

has ω -energy less than E

or is contained inside a d_ω -ball radius r ,

satisfies

$$|df|_\omega < c \text{ at } z = 0$$

We shall omit the proof of the above Lemma which is a standard bubbling argument, similar, but easier than the proof of the following:

Lemma 3.11. *Given $(\mathbf{B}, J, \Omega, \omega, g)$, for any $E > 0$,*

there exists

some $\epsilon > 0$,

distance $r > 0$,

and constant c ,

each depending continuously on $(\mathbf{B}, J, \Omega, \omega, g)$

so that any holomorphic map f of a disk $|z| \leq 1$ into $\mathbf{B}$ with Ω -energy less than E , and either

contained in a d_ω -ball of radius r

or having ω -energy less than ϵ

satisfies

$$|df|_g < c \text{ at } z = 0$$

Proof:

Suppose that this lemma was false. Then there would exist some sequence of maps f_i satisfying the above conditions with $|df_i(0)| \rightarrow \infty$. First, we obtain a sequence of rescaled J holomorphic maps $\tilde{f}_i : D(R_i) \rightarrow \mathbf{B}$ from the standard complex disk of radius R_i so that

$$\begin{aligned} |d\tilde{f}_i| &\leq 2 \\ |d\tilde{f}_i(0)| &= 1 \\ \lim_{i \rightarrow \infty} R_i &= \infty \end{aligned}$$

We achieve this in the same way as in any standard bubbling off argument such as the proof of lemma 5.11 in [1]. We can then use Lemma 3.7, Lemma 3.8, and Lemma 3.9 to get a bound on the higher derivatives of $\tilde{f}_i$ on $D(R_i - 1)$.

As $\mathbf{B}$ is complete, we can choose a subsequence so that $f_i(0)$ converges to some $p \in \mathbf{B}$. Lemma 3.1 tells us that the geometry of (g, J, ω) around $f_i(0)$ converges to that around p . Considering all our maps as maps sending 0 to p , we can choose a subsequence that converges on compact subsets to a non constant holomorphic map $f : \mathbb{C} \rightarrow \mathbf{B}$. Note that f either has ω -energy less than ϵ or is contained in a ball of radius r in the ω pseudo-metric. Because f must have finite ω energy, we can use Lemma 3.6 to tell us that f must converge in the ω pseudo-metric as $|z| \rightarrow \infty$. Then by choosing ϵ or r small enough, we can prove using Lemma 3.4 and the standard removable singularity theorem for holomorphic curves, that f must be constant in $[\mathbf{B}]$ because f either has ω energy less than ϵ or is contained in a d_ω -ball of radius r .

Now that we have that the image of f is a point in $[\mathbf{B}]$, we know that f must be contained entirely within some $(\mathbb{C}^*)^n$ worth of points over a single point in $[\mathbf{B}]$ and a single point in $\mathbf{B}$. This $(\mathbb{C}^*)^n$ has the standard complex structure, so f gives us n entire holomorphic maps from $\mathbb{C} \rightarrow \mathbb{C}^*$. As f is non constant, at least one of these maps must be non constant. This map must have infinite degree, as it must have dense image in the universal cover of $\mathbb{C}^*$ which is just the usual complex plane. It therefore has infinite Ω -energy, contradicting the fact that it must have Ω -energy less than E .

To prove the family case, it is important to note that (as stated at the start of this section), when we say our constants depend continuously on $(\mathbf{B}, J, \Omega, \omega, g)$, we mean that given any finite dimensional family, $(\hat{\mathbf{B}} \rightarrow \mathbf{G}, J, \Omega, \omega, g)$, the constants can be chosen to depend continuously on $\mathbf{G}$. With this understood, the proof in the family case is analogous to the above proof. $\square$

Lemma 3.12. *Given $(\mathbf{B}, J, \Omega, \omega)$,*

there exists

an energy bound $E > 0$,

a distance $r > 0$,

and for any $0 < \delta < 1$ a constant c ,

each depending continuously on $(\mathbf{B}, J, \Omega, \omega)$

so that any holomorphic map f from the annulus $e^{-(R+1)} < |z| < e^{(R+1)}$ to $\mathbf{B}$ with ω energy less than E , or contained inside a ball of ω radius r satisfies the following inequality

$$d_\omega(f(z), f(1)) \leq ce^{-\delta R}(|z|^\delta + |z|^{-\delta}) \text{ for } e^{-R} \leq |z| \leq e^R$$

Proof:

For this proof we shall use coordinates $z = e^{t+i\theta}$, and the cylindrical metric in which $\{\frac{\partial}{\partial t}, \frac{\partial}{\partial \theta}\}$ are an orthonormal frame.

We choose our energy bound $E > 0$ small enough that we can use the derivative bound from Lemma 3.10, and Lemma 3.6 implies that the smaller annulus is contained a small enough ball that we can use the coordinates of Lemma 3.7 (we also choose r small enough that this is true). We then have the following estimate for the (standard) $\bar{\partial}$ of f in these coordinates.

$$|\bar{\partial}f| \leq c_1 |df| |f|$$

We can run this through the inequality from Lemma 3.9 to obtain the following estimate on $\bar{\partial}f$ restricted to the interior of a disk (which holds on the interior of the cylinder where we have the derivative bound from Lemma 3.10.)

$$\|\bar{\partial}f\|_{L^p(D(\frac{1}{2}))} \leq c_1 \|f\|_{L^p_1(D(\frac{1}{2}))} \|f\|_{L^\infty(D(\frac{1}{2}))} \leq c_2 \|f\|_{L^\infty(D(1))} \|f\|_{L^\infty(D(\frac{1}{2}))} \quad (1)$$

The second inequality uses Lemma 3.9 and the derivative bound from Lemma 3.10. Here c_2 is a constant depending only on the p from L^p and $(\mathbf{B}, J, \omega)$. We can fix p to be something bigger than 2. By choosing E or r small, we can force $|f|$ to be as small as we like on the smaller cylinder using Lemma 3.6.

Now we can use Cauchy's integral formula

$$2\pi i f(z_0) = - \int_{|z|=1} \frac{f(z)}{z - z_0} dz + \int_{|z|=e^{2l}} \frac{f(z)}{z - z_0} dz + \int_{1 \leq |z| \leq e^{2l}} \frac{\bar{\partial}f(z)}{z - z_0} \wedge dz$$

or in our coordinates,

$$\begin{aligned} f(t_0, \theta_0) &= -\frac{1}{2\pi} \int_0^{2\pi} \frac{f(0, \theta)}{1 - e^{t_0} e^{i(\theta - \theta_0)}} d\theta + \frac{1}{2\pi} \int_0^{2\pi} \frac{f(2l, \theta)}{1 - e^{t_0 - 2l} e^{i(\theta - \theta_0)}} d\theta \\ &\quad + \frac{1}{2\pi} \int_0^{2\pi} \int_0^{2l} \frac{\bar{\partial}f(\theta, t)}{1 - e^{t_0 - t} e^{i(\theta - \theta_0)}} dt d\theta \end{aligned}$$

Let us now consider each term of this expression for $f(l, \theta_0)$ in the middle of a cylinder under the assumption that the average of $f(2l, \theta)$ is 0.

The first term:

$$\left| \frac{1}{2\pi} \int_0^{2\pi} \frac{f(0, \theta)}{1 - e^l e^{i(\theta - \theta_0)}} d\theta \right| \leq \frac{1}{e^l - 1} \max |f(0, \theta)|$$

The second term using that the average of $f(2l, \theta)$ is 0,

$$\begin{aligned} \left| \frac{1}{2\pi} \int_0^{2\pi} \frac{f(2l, \theta)}{1 - e^{l-2l} e^{i(\theta - \theta_0)}} d\theta \right| &= \left| \frac{1}{2\pi} \int_0^{2\pi} f(2l, \theta) \left(\frac{1}{1 - e^{-l} e^{i(\theta - \theta_0)}} - 1 \right) d\theta \right| \\ &= \left| \frac{1}{2\pi} \int_0^{2\pi} f(2l, \theta) \left(\frac{e^{-l} e^{i(\theta - \theta_0)}}{1 - e^{-l} e^{i(\theta - \theta_0)}} \right) d\theta \right| \\ &\leq \frac{1}{e^l - 1} \max |f(2l, \theta)| \end{aligned}$$

The third term:

$$\begin{aligned} \left| \frac{1}{2\pi} \int_0^{2\pi} \int_0^{2l} \frac{\bar{\partial} f(\theta, t)}{1 - e^{t_0 - t} e^{i(\theta - \theta_0)}} dt d\theta \right| &\leq \|\bar{\partial} f\|_{L^3} \frac{1}{2\pi} \left\| \frac{1}{1 - e^{l-t} e^{i(\theta - \theta_0)}} \right\|_{L^{\frac{3}{2}}} \\ &\leq c_3(l+1) \left(\max_{t \in [-1, 2l+1]} |f(t, \theta)| \right) \left(\max_{t \in [0, 2l]} |f(t, \theta)| \right) \end{aligned}$$

The constant c_3 depends only on $(\mathbf{B}, J, \omega)$. It is zero if J is integrable.

To summarize the above, we have the following expression which holds if the average of $f(\theta, 2l)$ is 0. (It also holds if the average of $f(\theta, 0)$ is 0 due to the symmetry of the cylinder.)

$$\begin{aligned} |f(l, \theta)| &\leq \frac{1}{e^l - 1} (\max |f(0, \theta)| + \max |f(2l, \theta)|) \\ &\quad + c_3(l+1) \left(\max_{t \in [-1, 2l+1]} |f(t, \theta)| \right) \left(\max_{t \in [0, 2l]} |f(t, \theta)| \right) \end{aligned} \quad (2)$$

The amount that the average of $f(t_0, \theta)$ changes with t_0 inside $[0, 2l]$ is determined by the integral of $\bar{\partial} f$, which is dominated as above by a term of the form

$$\text{change in average} \leq c_4 l \left(\max_{t \in [-1, 2l+1]} |f(t, \theta)| \right) \left(\max_{t \in [0, 2l]} |f(t, \theta)| \right) \quad (3)$$

Now let's put these estimates into slightly more invariant terms: Define the variation of $f(t, \theta)$ on $[a, b] \times \mathbb{T}^1$ as follows:

$$Vf([a, b]) := \max_{t_1, t_2 \in [a, b]} d_\omega(f(t_1, \theta_1), f(t_2, \theta_2))$$

Now the above two estimates give the following:

$$Vf([a, a+l]) \leq \left(\frac{8}{e^l - 1} + c_5(l+1) Vf([a-l-1, a+2l+1]) \right) Vf([a-l, a+2l]) \quad (4)$$

We shall now complete the argument for one particular δ - say $\delta = \frac{1}{2}$. Now if we choose l large enough, and then make Vf small enough by making our energy bound E small (or directly making r small), we can get the estimate

$$Vf([a, a+l]) \leq \frac{e^{-\delta l}}{3} Vf([a-l, a+2l]) \text{ for } [a-l-1, a+2l+1] \subset [-R, R] \quad (5)$$

Applying inequality 5 three times, we get that on the appropriate part of the cylinder,

$$\begin{aligned} Vf([a-l, a+2l]) &\leq e^{-\delta l} Vf([a-2l, a+3l]) \\ \text{so } Vf([a, a+l]) &\leq \frac{e^{-\delta 2l}}{3} Vf([a-2l, a+3l]) \end{aligned}$$

Inductively continuing this argument gives that if $[a-nl, a+(n+1)l] \subset [-R+1, R-1]$,

$$Vf([a, a+l]) \leq \frac{e^{-\delta nl}}{3} Vf([a-nl, a+(n+1)l])$$

The required estimate follows from this.

Once we have the required convergence for one δ , we may obtain inequality 5 for other δ from 4 by restricting suitably far into the interior of the annulus so that Vf is small. Then repeating the above argument gives the required inequality (with a different constant). $\square$

Lemma 3.9, Lemma 3.8, and the coordinates from Lemma 3.7 from page 11 can be used with inequality 1 on page 14 in a standard fashion to get a similar estimate on the derivative of f in the ω pseudo metric. This gives the following corollary which states with quantifiers that any cylinder with small enough ω -energy can be restricted to a smaller cylinder to have ω -energy as small as you like.

Corollary 3.13. *Given $(\mathbf{B}, J, \Omega, \omega)$,*

there exists some energy $E > 0$ depending continuously on $(\mathbf{B}, J, \Omega, \omega)$

so that given any $E' > 0$,

there exists some distance R depending continuously on $(\mathbf{B}, J, \Omega, \omega, E')$

so that given any holomorphic map f of a cylinder

$$\{e^{-l-R} < |z| < e^{l+R}\}$$

to $\mathbf{B}$ with ω energy less than E ,

the ω energy of f restricted to $e^{-l} < |z| < e^l$ is less than E' .

Lemma 3.14. *Given $(\mathbf{B}, J, \Omega, g)$, an energy bound $E < \infty$, and a covering of $\mathbf{B}$ by coordinate charts,*

there exists

an open cover $\{U_i\}$ of $\mathbf{B}$

and for each $0 < \delta < 1$ a constant c

so that the following is true:

Each member of our open cover U_i is contained in some particular coordinate chart.

Given any holomorphic map f with Ω -energy less than E from a cylinder $e^{-(R+1)} \leq |z| \leq e^{(R+1)}$ to $\mathbf{B}$ contained inside U_i ,

there exists a map F given in coordinates as

$$F(z) := (c_1 z^{\alpha_1} \mathbf{t}^{a_1}, \dots, c_k z^{\alpha_k} \mathbf{t}^{a_k}, c_{k+1}, \dots, c_n)$$

so that

$$\text{dist}(f(z) - F(z)) \leq ce^{-\delta R} \left(|z|^\delta + |z|^{-\delta} \right) \text{ for } e^{-R} \leq |z| \leq e^R$$

where for $i \in [1, k]$, $c_i \in \mathbb{C}^$, $a_i \in \mathbb{R}$, $\alpha_i \in \mathbb{Z}$, and for $i \in [k+1, n]$, $c_i \in \mathbb{R}$. dist indicates distance in the metric g .*

In the case that we have a family $(\hat{\mathbf{B}} \longrightarrow \mathbf{G}, J, \Omega, g)$ with the above structure, we can choose c continuous on $\mathbf{G}$.

Proof:

First, choose any symplectic form $\omega \in \Omega$. Because $\mathbf{B}$ is complete, we can then choose a finite covering $\{U_i\}$ of $\mathbf{B}$ by open subsets which are contained inside our coordinate charts and which are small enough in the ω pseudo metric so that Lemma 3.12 tells us that for any holomorphic map from a cylinder as above contained inside U_i ,

$$d_\omega(f(z), f(1)) \leq ce^{-\delta'R}(|z|^{\delta'} + |z|^{-\delta'}) \quad (6)$$

where $\delta' = \frac{\delta+1}{2}$. In the family case, we choose our open cover $\{U_i\}$ to be fiberwise finite, and the above constant c depends continuously on $\mathbf{G}$.

In our coordinates,

$$f(1) = (c_1 \mathbf{t}^{a_1}, \dots, c_k \mathbf{t}^{a_k}, c_{k+1}, \dots, c_n)$$

We can choose α_i so that the winding numbers of the first k coordinates of $f(e^{i\theta})$ are the same as our model map F :

$$F := (c_1 z^{\alpha_1} \mathbf{t}^{a_1}, \dots, c_k z^{\alpha_k} \mathbf{t}^{a_k}, c_{k+1}, \dots, c_n)$$

The metric g can be compared to the pseudo metric from ω on the last coordinates $c_{k+1}, \dots, c_n$, so we only need to prove convergence in the first k coordinates. Let us do this for f restricted to the first coordinate f_1 .

For this, use coordinates $z = e^{t+i\theta}$ on the domain with the usual cylindrical metric and the similar cylindrical metric on our target. (Choose members of our open cover to be compactly contained inside our coordinate charts so that this standard flat metric on the coordinate chart is comparable to g). Use the notation

$$\frac{f_1(z)}{c_1 z^{\alpha_1} \mathbf{t}^{a_1}} = e^{h(z)}$$

where h is a $\mathbb{C}$ valued function so that $h(0) = 0$. The metric we are using on the target is the standard Euclidean metric on $\mathbb{C}$. Our goal is to prove the appropriate estimate for h .

First note that Lemma 3.9 and Lemma 3.8 can be used with the inequality 6 in a standard fashion to get a similar estimate on the derivative of f in the ω pseudo metric.

$$|df(z)|_\omega \leq ce^{-\delta'R}(|z|^{\delta'} + |z|^{-\delta'}) \text{ for } e^{-R} \leq |z| \leq e^R \quad (7)$$

(The c in the above inequality is some new constant which is independent of f). So long as we have chosen members of our open cover U_i compactly contained inside our coordinate charts, the following inequality holds inside U_i :

$$|Jv - J_0 v|_g \leq M|v|_\omega$$

where J_0 indicates the standard complex structure, and M is a constant, which may have to depend on $\mathbf{G}$ in the family case. To see this, consider the standard basis for vectorfields in coordinates $\frac{\partial}{\partial x_i}$ and the real and imaginary parts of $\tilde{z}_i \frac{\partial}{\partial \tilde{z}_i}$, and suppose that v is one of these basis vector fields. $Jv - J_0 v$ is smooth

and vanishes whenever $|v|_\omega = 0$. The fact that d_ω is actually a metric on the smooth part of our coordinate chart then implies that the above inequality holds for v being a basis vector field. As each side of the above inequality measures the size of a tensor, the general case follows. The above inequality implies that $|\bar{\partial}f_1|_g$ in our coordinates is controlled by $|df|_\omega$. This then gives that for some new constant c independent of f we get the following inequality:

$$|\bar{\partial}h(z)| \leq ce^{-\delta'R}(|z|^{\delta'} + |z|^{-\delta'}) \text{ for } e^{-R} \leq |z| \leq e^R \quad (8)$$

If we choose our open cover small enough using d_ω , Lemma 3.11 gives us a bound for $|df|_g$ on the interior of the cylinder, so we have that there exists some c independent of f so that

$$|dh(z)| < c \text{ for } e^{-R} \leq |z| \leq e^R \quad (9)$$

We can now proceed roughly as we did in the proof of Lemma 3.12. In particular, Cauchy's integral theorem tells us that

$$\begin{aligned} h(t_0, \theta_0) &= -\frac{1}{2\pi} \int_0^{2\pi} \frac{h(t_0 - l, \theta)}{1 - e^l e^{i(\theta - \theta_0)}} d\theta + \frac{1}{2\pi} \int_0^{2\pi} \frac{h(t_0 + l, \theta)}{1 - e^{-l} e^{i(\theta - \theta_0)}} d\theta \\ &\quad + \frac{1}{2\pi} \int_0^{2\pi} \int_{t_0-l}^{t_0+l} \frac{\bar{\partial}h(t, \theta)}{1 - e^{t_0-t} e^{i(\theta - \theta_0)}} dt d\theta \end{aligned}$$

We can now bound each term in the above expression as in the proof of Lemma 3.12, except we use the estimate 8 to bound $|\bar{\partial}h|$.

Then we have the following estimate:

$$\begin{aligned} \left| h(t_0, \theta_0) - \frac{1}{2\pi} \int_0^{2\pi} h(t_0 + l, \theta) d\theta \right| &\leq \frac{1}{e^l - 1} \max \left| h(t_0 - l, \theta) - \frac{1}{2\pi} \int_0^{2\pi} h(t_0 + l, \theta) d\theta \right| \\ &\quad + \frac{1}{e^l - l} \max \left| h(t_0 + l, \theta) - \frac{1}{2\pi} \int_0^{2\pi} h(t_0 + l, \theta) d\theta \right| \\ &\quad + c(l+1)e^{-\delta'R}e^{\delta'l} \left(e^{\delta't_0} + e^{-\delta't_0} \right) \end{aligned}$$

(Of course, this is a new constant c , which is independent of l or h .)

Note that the change in the average of h is determined by the integral of $\bar{\partial}h$, which we can bound using estimate 8. The change in this average can then be absorbed into the last term of the above inequality. Define the variation of h for a particular t as follows:

$$Vh(t) := \max_{\theta} \left| h(t, \theta) - \frac{1}{2\pi} \int_0^{2\pi} h(t, \theta) d\theta \right|$$

We then have the following estimate (with a new constant c):

$$Vh(t) \leq \frac{1}{e^l - 1} (Vh(t+l) + Vh(t-l)) + c(l+1)e^{\delta'(l-R)} \left(e^{\delta't} + e^{-\delta't} \right)$$

Recalling that $\delta < \delta'$ and Vh is bounded by equation 9, we can choose l large enough so that for R sufficiently large, using the above estimate recursively tells us that there exists some c independent of f or R so that

$$Vh(t) \leq ce^{-\delta R}(e^{\delta t} + e^{-\delta t}) \text{ for } -R \leq t \leq R$$

(This estimate follows automatically from the bound on Vh for R bounded.) The required estimate for h then follows from the fact that the change in the average of h is bounded by the estimate 8, which is stronger than what we need (which is the same equation with δ in place of δ'). We therefore have

$$h(z) \leq ce^{-\delta R}(|z|^\delta + |z|^{-\delta}) \text{ for } e^{-R} \leq |z| \leq e^R$$

which is the required estimate. $\square$

To prove compactness results, we shall divide our domain up into annuli with small energy, and other compact pieces with derivative bounds. For this we shall need some facts about annuli. Recall the following standard definition for the conformal modulus of a Riemann surface which is an annulus:

Definition 3.15. *The conformal modulus of an annulus A is defined as follows. Let $S(A)$ denote the set of all continuous functions with L^2 integrable derivatives on A which approach 1 at one boundary of A and 0 at the other. Then the conformal modulus of A is defined as*

$$R(A) := \sup_{f \in S(A)} \frac{2\pi}{\int_A (df \circ j) \wedge df} = \sup_{f \in S(A)} \frac{2\pi}{\int_A |df|^2}$$

We can extend the definition of conformal modulus to include ‘long’ annuli inside exploded curves as follows:

Definition 3.16. *Call a (non complete) exploded curve $\mathbf{A}$ an exploded annulus of conformal modulus $\log x\mathbf{t}^{-l}$ if it is connected, and there exists an injective holomorphic map $f : \mathbf{A} \rightarrow \mathbf{T}$ with image $\{1 < |\tilde{z}| < x\mathbf{t}^{-l}\}$. Call it an exploded annulus with semi infinite conformal modulus if it is equal to (a refinement of) $\{|\tilde{z}| < 1\} \subset \mathbf{T}_1^1$.*

(We use $x\mathbf{t}^{-l}$ in the above definition because the tropical part of the resulting annulus will have length l .) Two exploded annuli with the same conformal modulus may not be isomorphic, but they will have a common refinement.

We shall need the following lemma containing some useful properties of the (usual) conformal modulus.

Lemma 3.17. *1. An open annulus A is conformally equivalent to*

$$\{1 < |z| < e^R\}$$

if and only if the conformal modulus of A is R . If the conformal modulus of A is infinite, then A is conformally equivalent to either a punctured disk, or a twice punctured sphere.

2. If $\{A_i\}$ is a set of disjoint annuli $A_i \subset A$ none of which bound a disk in A , then

$$R(A) \geq \sum R(A_i)$$

3. If A_1 and A_2 are annuli contained inside the same Riemann surface which share a boundary so that


(a) $R(A_2) < \infty$,

(b) and the other boundary of A_1 intersects the other boundary of A_2 and the circle at the center of A_2 ,

then


$$R(A_2) + \frac{16\pi^2}{R(A_2)} \geq R(A_1)$$

(This inequality is not sharp.)



4. If A_1 and A_2 are annuli contained inside a Riemann surface so that every circle homotopic to the boundary inside A_2 contains a segment inside A_1 that intersects both boundaries of A_1 , then

$$R(A_1) \leq \frac{4\pi^2}{R(A_2)}$$



Proof:

The first two items are well known. To prove item 3, first set $R = R(A_2)$ and put coordinates $(0, R) \times \mathbb{R}/2\pi\mathbb{Z}$ on A_2 . Consider any function $f \in S(A_1)$. Without losing generality, we can assume that the shared boundary is $(0, \theta)$, and that some segment of the other boundary of A_1 is a curve in A_2 between $(\frac{R}{2}, 0)$ and (R, θ_0) where $\theta_0 \geq 0$. Then consider integrating $|df|^2$ along diagonal lines $(Rt, -4\pi t + c)$. Each of these lines contains a segment inside A_1 on which the integral of df is 1. The length of each segment is bounded by $((4\pi)^2 + R^2)^{\frac{1}{2}}$. The integral of $|df|^2$ along this segment is therefore at least $(16\pi^2 + R^2)^{-\frac{1}{2}}$. This tells us that the integral of $|df|^2$ over A_1 is at least $\frac{2\pi R}{((4\pi)^2 + R^2)}$, and therefore,

$$R(A_1) \leq \frac{(16\pi^2 + R^2)}{R} = R(A_2) + \frac{16\pi^2}{R(A_2)}$$

To prove item 4, consider a function $f \in S(A_1)$. We shall integrate $|df|^2$ along segments of the form (c, t) inside $A_2 \cap A_1$ traveling from one boundary of A_1 to the other. The integral of df along such a segment is 1, and the length of the segment is at most 2π , so the integral of $|df|^2$ along the segment is at least $\frac{1}{2\pi}$, and the integral of $|df|^2$ over A_1 is at least $\frac{R(A_2)}{2\pi}$. Therefore, we have

$$R(A_1) \leq \frac{4\pi^2}{R(A_2)}$$

□

The following proposition gives the existence of a decomposition of any holomorphic curve f with bounded energy and topology. This decomposition is into annular regions with small energy and complementary regions with bounded conformal modulus on which the derivative of f is bounded.

Proposition 3.18.

Given

$(\mathbf{B}, J, \Omega, \omega, g),$

an energy bound E ,

a number N to bound genus and number of punctures,

and small enough $\epsilon > 0$ (which will bound the energy of some annular regions),

there exists

*a number bound M depending lower semicontinuously on $(\mathbf{B}, J, \Omega, \omega, g, E, N, \epsilon)$
(M shall bound a number of annular regions)*

for any large enough distance R , a derivative bound c and conformal bound $\tilde{R}$ depending continuously on $(\mathbf{B}, J, \Omega, \omega, g, E, N, \epsilon, R)$

and an energy lower bound $\epsilon_0 > 0$ depending continuously on (E, R, ϵ)

so that the following is true:

Given any complete, stable holomorphic curve

$$f : \mathbf{C} \longrightarrow \mathbf{B}$$

with energy at most E , and with genus and number of punctures at most N , there exists some collection of at most M exploded annuli $\mathbf{A}_i \subset \mathbf{C}$, so that:

1. *Each $\mathbf{A}_i$ has conformal modulus larger than $2R$*
2. *Put the standard cylindrical metric on $\mathbf{A}_i$, and use the notation $\mathbf{A}_{l,i}$ to denote the annulus consisting of all points in $\mathbf{A}_i$ with distance to the boundary at least l . Then*

$$\mathbf{A}_{\frac{R}{2},i} \cap \mathbf{A}_{\frac{R}{2},j} = \emptyset \text{ if } i \neq j$$

3. *f restricted to $\mathbf{A}_i$ has energy less than ϵ .*


4. Each component of $\mathbf{C} - \bigcup \mathbf{A}_{R,i}$ is a smooth Riemann surface with bounded conformal geometry in the sense that any annulus inside one of these smooth components with conformal modulus greater than $\hat{R}$ must bound a smooth disk inside that component.
5. The following metrics can be put on $\mathbf{A}_i$ and each smooth component of $\mathbf{C} - \bigcup \mathbf{A}_{R,i}$:
 - (a) On any component which is a smooth torus, use the unique flat metric in the conformal class of the complex structure so that the area of the torus is 1.
 - (b) If the component is equal to a disk, an identification with the standard unit disk can be chosen so that if $\mathbf{A}_{R,i}$ is the bounding annulus, 0 is in the complement of $\mathbf{A}_i$. Give components such as this the standard Euclidean metric.
 - (c) If the component is equal to some annulus, give it the standard cylindrical metric on $\mathbb{R}/\mathbb{Z} \times (0, l)$. Give each $\mathbf{A}_i$ the analogous standard metric.
 - (d) Any component not equal to a torus, annulus, or disk will admit a unique metric in the correct conformal class with curvature -1 so that boundary components are geodesic (there will be no components which are smooth spheres). Give these components this metric.

On any component of $\mathbf{C} - \bigcup \mathbf{A}_{\frac{9R}{10},i}$, the derivative in this metric is bounded by c

$$|df|_g < c$$

Moreover, on $\mathbf{A}_{\frac{6R}{10},i} - \mathbf{A}_{\frac{9R}{10},i}$, the ratio between the two metrics defined above is less than c .

6. f restricted to any component of $\mathbf{C} - \bigcup \mathbf{A}_{R,i}$ which is a disk, annulus or torus has ω energy greater than $\epsilon_0 > 0$.




Proof:

Lemma 3.11 tells us that for ϵ small enough, any holomorphic map of the unit disk with ω -energy less than ϵ and Ω -energy less than E must have derivative at 0 bounded by c_0 . We shall prove our theorem for ϵ small enough so that this is true, and small enough that Corollary 3.13 also holds with an energy bound of ϵ . We shall also choose our distance R greater than 4π (for use with Lemma 3.17), and large enough that Corollary 3.13 tells us that if f restricted to some smooth annulus $\mathbf{A}_i$ has ω -energy less than ϵ , then the ω -energy of f restricted to the smaller annulus $\mathbf{A}_{\frac{R}{2},i}$ is less than $\frac{\epsilon}{5}$.


Let us now begin to construct our annuli. First, note that any edge of $\mathbf{C}$ has zero ω -energy, so we can choose an exploded annulus $\mathbf{A}_i$ containing each edge with ω -energy less than $\frac{\epsilon}{5}$. (Note for use with item 6 that in the case that this bounds a disk, the ω -energy of the resulting disk will be at least $\frac{4\epsilon}{5}$.) We can do this so that these $\mathbf{A}_i$ are mutually disjoint. Note that the complement of these $\mathbf{A}_i$ is a smooth Riemann surface with boundary. (Note also that no component of $\mathbf{C}$ can be isomorphic to $\mathbf{T}$ as the Ω -energy of any complete component is equal to the ω -energy which must be zero on $\mathbf{T}$.)

For each connected component of $\mathbf{C}$ which is a smooth sphere, we shall now remove an annulus. First, note that we can put some round metric in the correct conformal class of the sphere so that there exist 3 mutually perpendicular geodesics which divide our sphere into 8 regions each of which has equal ω -energy. As our sphere must have energy at least ϵ in order to be stable, we can choose two antipodal regions that each have energy $\frac{\epsilon}{8}$. By choosing $\epsilon_0 > 0$ small enough, we can then get that there exist two disks with ω -energy at least ϵ_0 which intersect each of these regions, and which have radius as small as we like. Choose $\epsilon_0 > 0$ small enough that the complement of these disks is an annulus of conformal radius at least $k(2R + 1)$ for some integer $k > \frac{5E}{\epsilon}$. How small ϵ_0 is required to achieve this depends only on R , ϵ and E . Then we can divide this annulus into k annuli with conformal modulus at least $(2R + 1)$, at least one of

which has ω -energy at most $\frac{\epsilon}{5}$. Add this annulus to our collection. Note that it bounds disks which have energy at least ϵ_0 .




Consider a holomorphic injection of the unit disk $i : D \rightarrow \mathbf{C}$ into the complement of our annuli $\mathbf{A}_{R,i}$ constructed up to this point. Suppose that $i(0)$ is in the complement of $\mathbf{A}_{\frac{R}{4},i}$. Then Lemma 3.17 tells us that the restriction of i to $|z| < e^{-\frac{16\pi^2}{R}}$ is in the complement of $\mathbf{A}_{\frac{R}{2},i}$. Now suppose that $|d(f \circ i)| > c_0 e^{\frac{16\pi^2}{R} + k(2R+1)}$ where k is some integer greater than $\frac{5E}{\epsilon}$. Then the restriction of f to the disk $|z| < e^{-\frac{16\pi^2}{R} - k(2R+1)}$ must have energy greater than ϵ due to Lemma 3.11. This disk is surrounded by k disjoint annuli of conformal modulus $(2R+1)$ which are contained in the complement of $\mathbf{A}_{\frac{R}{2},i}$. At least one of these must have energy less than $\frac{\epsilon}{5}$. Add this annulus to our collection. Continue adding annuli in this manner. After a finite (but not universally bounded) number of times, no more annuli can be added in this way. We shall argue this below after adding some more annuli.



We now want to add annuli so that condition 4 is satisfied. Suppose some smooth component of $\mathbf{C} - \bigcup \mathbf{A}_{R,i}$ contains some annulus of conformal modulus greater than $2R(3 + \frac{5E}{\epsilon})$ that does not bound a disk. Note that as we have chosen


$R > 4\pi$, the size of such an annulus is greater than $2R(1 + \frac{5E}{\epsilon}) + 2R + \frac{32\pi^2}{R}$. The annulus of size R at its end can't be contained entirely within $\mathbf{A}_i$, and then applying Lemma 3.17 part 4 with the next annulus of size $\frac{16\pi^2}{R}$ in the place of A_1 , we see that our annulus minus annuli at each end of conformal modulus $R + \frac{16\pi^2}{R}$ must be contained entirely in the complement of $\mathbf{A}_{\frac{R}{2},i}$. We then obtain more than $\frac{5E}{\epsilon}$ disjoint annuli in the complement of $\mathbf{A}_{\frac{R}{2},i}$ with conformal modulus greater than $2R$. The restriction of f to at least one of these annuli must have ω -energy less than ϵ . Add this annulus to our collection.



We shall now argue that we can only add a finite number of annuli in this manner. The number of annuli that bound disks is bounded by $\frac{E}{\epsilon}$ plus the twice the number of connected components of $\mathbf{C}$ that are spheres (which is bounded by $\frac{E}{\epsilon}$). There are also a finite number of exploded annuli with infinite conformal modulus. The number of these is bounded by our topological bounds, and the fact that there are at most $\frac{E}{\epsilon}$ spherical components with fewer than 3 punctures. Call the complement of all the above annuli $\mathbf{C}_0$. There is then a bound on the number of homotopy classes in $\mathbf{C}_0$ of our remaining annuli, given by our bounds on the topology of $\mathbf{C}$, the fact that we have removed a bounded number of annuli, and the observation that for any Riemann surface with boundary, there are only a finite number of homotopy classes which contain annuli of conformal modulus greater than 4π (Lemma 3.17 part 4 can help to prove this). If there were an infinite number of annuli $\mathbf{A}_{\frac{R}{2},i}$ in our collection in the same homotopy class, then there would exist an annulus in $\mathbf{C}_0$ in that homotopy class which contains all of them. As the $\mathbf{A}_{\frac{R}{2},i}$ are disjoint and have conformal modulus at least R , this annulus would have to have infinite conformal modulus. Note that there are no nontrivial annuli of infinite conformal modulus in $\mathbf{C}_0$ (because we have removed the annuli containing edges and at least one annulus from each spherical component), and if an annulus of infinite conformal modulus surrounds a disk with energy greater than ϵ_0 , then the derivative of f must have been unbounded there. This is not possible, so we must only have a finite number of annuli in our collection. Note that everything apart from our bound on the number of annuli in a fixed homotopy class is bounded independent of f .

Now we shall merge some annuli so that there exists no annular component of $\mathbf{C} - \bigcup \mathbf{A}_{R,i}$ with ω -energy less than $\frac{\epsilon}{5}$. Then we will have a bound on the number of annuli which is independent of f . Suppose that we have some collection $\{\mathbf{A}_1, \dots, \mathbf{A}_n\}$ of our annuli so that $\mathbf{A}_{R,i}$ and $\mathbf{A}_{R,i+1}$ bound an annulus which has ω -energy less than $\frac{\epsilon}{5}$. Use the notation $\mathbf{A}_{[m,n]}$ to denote the annulus that consists of $\mathbf{A}_m$, $\mathbf{A}_n$ and everything in between. Then, as we've chosen each of our $\mathbf{A}_i$ to have ω -energy less than $\frac{\epsilon}{5}$, $\mathbf{A}_{[i,i+2]}$ has ω -energy less than ϵ , so we can apply Corollary 3.13 to show that far enough into the interior of

this cylinder, there is very little ω -energy. (Note that as the edges of exploded curves always have no ω -energy, so there is no difficulty in applying this lemma in the seemingly more general setting of exploded annuli.) Applying Lemma 3.17 part 3, and noting that as we have chosen $R > 4\pi$, we have $2R > R + \frac{16\pi^2}{R}$, we see that $\mathbf{A}_{[i,i+2]} - \mathbf{A}_i - \mathbf{A}_{i+2} \subset \mathbf{A}_{\frac{R}{2},[i,i+2]}$. We have chosen R large enough that Corollary 3.13 tells us that the energy of f restricted to $\mathbf{A}_{\frac{R}{2},[i,i+2]}$ is less than $\frac{\epsilon}{5}$. We can now repeat this argument inductively to show that the energy of f restricted to $\mathbf{A}_{[1,n]}$ is less than ϵ and $\mathbf{A}_{\frac{R}{2},[1,n]}$ is less than $\frac{\epsilon}{5}$.



Now, replace all of our sets of annuli $\{\mathbf{A}_1, \dots, \mathbf{A}_n\}$ of maximal size obeying the above conditions with the annulus given by $\mathbf{A}_{\frac{R}{2},1}$, $\mathbf{A}_{\frac{R}{2},n}$ and everything in between. (Note that if we had some collection of annuli which bounded small energy annuli in a cyclic fashion, we would obtain a connected component of $\mathbf{C}$ which had energy less than ϵ , and was therefore not stable.)

We shall now check that all the conditions we require are satisfied by this new set of annuli. First, the number is bounded independent of f . The size of each annulus is greater than $2R$. Our resulting set of annuli also obey the non-intersection condition 2 because the original set did. We showed above that the energy of each new annulus is less than ϵ . Because the complement of our original set of annuli obeyed condition 4 with a bound of $2R(3 + \frac{5E}{\epsilon})$, by increasing our conformal bound appropriately and using Lemma 3.17, we can achieve condition 4. The lower bound on the ω -energy of unstable components is also satisfied by construction.

All that remains is condition 5. Note first that the complement of our annuli $\mathbf{A}_{R,i}$ admits metrics as described in 5. The fact that there exists some c so that the metrics we choose on different components differ by a factor of less than c on $\mathbf{A}_{\frac{6R}{10},i} - \mathbf{A}_{\frac{9R}{10},i}$ follows from conditions 4 and 2, and the bound on topology as follows: The complement of $\mathbf{A}_{R,i}$ is a Riemann surface with bounded genus, a bounded number of boundary components and bounded conformal geometry from condition 4. An easy compactness argument tells us that there exists a constant c' so that any injective holomorphic map of the unit disk into either $\mathbf{A}_i$ or the complement of $\mathbf{A}_{R,i}$ has derivative at 0 bounded by c' . We can use this (remembering that $R > 4\pi > 10$ so we can get a disk of unit size centered on any point inside $\mathbf{A}_{\frac{6R}{10}} - \mathbf{A}_{\frac{9R}{10}}$) to get the ratio of the metric on the complement of $\mathbf{A}_{R,i}$ divided by the metric on $\mathbf{A}_i$ is bounded by c' here.

Then, we can use Lemma 3.17 and our conformal bound from condition 4 to get some lower bound $r_0 > 0$ so that the boundary of $\mathbf{A}_{\frac{9R}{10}}$ and $\mathbf{A}_{\frac{6R}{10}}$ is further than r_0 from the boundary of the complement of $\mathbf{A}_{R,i}$. A second easy compactness argument then tells us that if we fix any distance $r_0 > 0$, there exists some constant c_{r_0} (only depending on the bounds in the above paragraph) so that for every point further than r_0 from the boundary there exists an injective holomorphic map of the unit disk sending 0 to that point, with derivative at 0

greater than c_{r_0} . We then get that the above ratio of metrics above is bounded below by $\frac{c_{r_0}}{c'}$.

We must now prove our derivative bound on the complement of $\mathbf{A}_{\frac{9R}{10}}$. We shall do this by proving that given any holomorphic injection of the unit disk $\iota : D \rightarrow \mathbf{C} - \bigcup \mathbf{A}_{R,i}$ so that $\iota(0) \notin \mathbf{A}_{\frac{9R}{10},i}$, then $|d(f \circ \iota)|$ is bounded. Then, our conformal bound from condition 4 will tell us that we have a bound on $|df|$ in the metric that we have chosen.

We do this in two cases. First, suppose that $\iota(0)$ is in the complement of $\mathbf{A}_{\frac{R}{4},i}$ for all our old annuli. Then, Lemma 3.17 tells us that the image of ι restricted to the disk of radius $e^{-16\frac{\pi^2}{R}}$ is contained in the complement $\mathbf{A}_{\frac{R}{2},i}$ for all our old annuli, as argued above. If the derivative of $f \circ \iota$ on this disk was large enough, we would be able to add another annulus to our old collection in the manner described above. As this process terminated, $|d(f \circ \iota)|$ is bounded. Second, suppose that $\iota(0)$ is contained inside $\mathbf{A}_{\frac{R}{4},i}$ for one of our old annuli. Then Lemma 3.17 tells us that the restriction of ι to the disk of radius $e^{-16\frac{\pi^2}{R}}$ must be contained inside $\mathbf{A}_i$, and therefore have energy less than ϵ . This means that $|d(f \circ \iota)| \leq c_0 e^{16\frac{\pi^2}{R}}$, and our derivative is bounded as required. $\square$

4 Compactness

Theorem 4.1. *Given a basic, complete exploded manifold $\mathbf{B}$ with a civilized almost complex structure J and strict taming Ω , the moduli stack $\mathcal{M}_{g,n,E}(\mathbf{B})$ of stable holomorphic curves with a fixed genus g and number of punctures n , and with Ω energy less than E is compact using the $C^{\infty,1}$ topology.*

More generally, if $(\hat{\mathbf{B}} \rightarrow \mathbf{G}, J, \Omega)$ is a family of such $(\mathbf{B}, J, \Omega)$, the map $\mathcal{M}_{g,n,E}(\hat{\mathbf{B}} \rightarrow \mathbf{G}) \rightarrow \underline{\mathbf{G}}$ is proper.

In particular, this means that given any sequence of the above holomorphic curves in the fibers over a convergent sequence of points in $\mathbf{G}$, there exists a subsequence f^i which converges to a holomorphic curve f as follows: There exists a sequence of families of $C^{\infty,1}$ curves,

$$\begin{array}{ccc} (\hat{\mathbf{C}}, j_i) & \xrightarrow{\hat{f}^i} & \hat{\mathbf{B}} \\ \downarrow & & \downarrow \\ \mathbf{F} & \longrightarrow & \mathbf{G} \end{array}$$

so that this sequence of families converges in $C^{\infty,1}$ to the family

$$\begin{array}{ccc} (\hat{\mathbf{C}}, j) & \xrightarrow{\hat{f}} & \hat{\mathbf{B}} \\ \downarrow & & \downarrow \\ \mathbf{F} & \longrightarrow & \mathbf{G} \end{array}$$

and a sequence of points p^i in $\mathbf{F}$ so that $p^i \rightarrow p$, f^i is the map given by the restriction of $\hat{f}^i$ to the fiber over p^i , and f is given by the restriction of $\hat{f}$ to the fiber over p . Of course, the case where we just have a single $(\mathbf{B}, J, \omega)$ is the same as the family case when $\mathbf{G}$ is a point.

The proof of this theorem uses Lemma 3.14 on page 16 and Proposition 3.18 on page 21. Together, these allow us to decompose holomorphic curves into pieces with bounded behavior. A standard Arzela-Ascoli type argument gets a type of convergence of these pieces. If we were working in the category of smooth manifolds, this would be sufficient to prove the compactness theorem because the type of convergence involved would have a unique limit, and therefore the limiting pieces would glue together. The extra problem that must be dealt with in this case is that we are not dealing with a type of convergence that has a unique limit, so we must work much harder to show that our resulting ‘limiting family’ for each piece glues together to the limiting family $\mathbf{F}$.

Example 4.2.

The following are examples of the types of non uniqueness we have to deal with:

1. In $\mathbf{T}_1^1$, let p^i be a sequence of points so that $\tilde{z}(p^i) = \frac{1}{i}t^0$. Then $p^i \rightarrow p$ where $p \in \mathbf{T}_1^1$ is any point so that $\tilde{z}(p) = ct^a$ where $a > 0$. Note that there is no ‘point’ $p \in \mathbf{T}_1^1$ so that $\tilde{z}(p) = 0t^0$. This is one reason that non unique limits need to be considered.
2. Expanding example 1, consider the maps $f^i : \mathbb{C}P^1 \rightarrow \mathbb{C}P^1 \times \mathbf{T}_1^1$ given by $f^i(u) = (u, p^i)$. This sequence of maps does not have a unique limit. We instead consider the family

$$\begin{array}{ccc} \mathbb{C}P^1 \times \mathbf{T}_1^1 & \xrightarrow{\hat{f} := \text{id}} & \mathbb{C}P^1 \times \mathbf{T}_1^1 \\ \downarrow & & \\ \mathbf{T}_1^1 & & \end{array}$$

Then the sequence of maps f^i are given by the restriction of $\hat{f}$ to the inverse image of p^i . These converge to $\hat{f}$ restricted to the inverse image of p , where $p \in \mathbf{T}_1^1$ is any point so that $p^i \rightarrow p$.

3. Expanding example 1 to see different behavior, consider the map

$$\pi : \mathbf{T}_2^2 \rightarrow \mathbf{T}_1^1$$

so that $\pi(\tilde{w}_1, \tilde{w}_2) = \tilde{w}_1\tilde{w}_2$. This can be considered to be a family of annuli by restricting to the subset where $|\tilde{w}_1| < 1$ and $|\tilde{w}_2| < 1$. Then $\pi^{-1}(p^i)$ is an annulus $1 > |\tilde{w}_1| > \frac{1}{i}$, or $\frac{1}{i} < |\tilde{w}_2| < 1$. These domains converge to $\pi^{-1}(p)$ where $p = ct^a$ for any $c \in \mathbb{C}^*$ and $a > 0$. $\pi^{-1}(ct^a)$ is equal to the subset of $\mathbf{T}_{[0,a]}^1$, with coordinate $\tilde{w}_1$ so that $1 > |\tilde{w}_1| > ct^a$

4. Let us expand on example 3 by adding in the information of maps f^i from our domains $\pi^{-1}(p^i)$ to $\mathbf{T}^2$. Suppose that

$$f^i(\tilde{w}_1) = (\tilde{c}_1^i \tilde{w}_1, \tilde{c}_2^i \tilde{w}_1)$$

so of course,

$$f^i(\tilde{w}_2) = (\tilde{c}_1^i \frac{1}{i} \tilde{w}_2^{-1}, \tilde{c}_2^i \frac{1}{i} \tilde{w}_2^{-1})$$

If $(c_1^i, c_2^i) \in \mathbf{T}^2$ are generically chosen, we will not have a unique limit even if we restrict to the bounded domain $1 > |\tilde{w}_1| > \frac{1}{2}$. To put all our f^i into an individual family, consider the family

$$\begin{array}{ccc} \mathbf{T}^2 \times \mathbf{T}_1^1 & \xrightarrow{\hat{f}(\tilde{c}_1, \tilde{c}_2, \tilde{w}_1, \tilde{w}_2) := (\tilde{c}_1 \tilde{w}_1, \tilde{c}_2 \tilde{w}_1)} & \mathbf{T}^2 \\ \downarrow \text{id} \times \pi & & \\ \mathbf{T}^2 \times \mathbf{T}_1^1 & & \end{array}$$

Our individual maps f^i are the restriction of $\hat{f}$ to the fiber over $(\tilde{c}_1^i, \tilde{c}_2^i, p^i)$. These converge to the restriction of $\hat{f}$ to the fiber over any point $(\tilde{c}_1, \tilde{c}_2, ct^a)$ where $a > 0$. Note that if we restricted to the domains where $1 > |\tilde{w}_1| > |\tilde{w}_2|$ or $1 > |\tilde{w}_2| > |\tilde{w}_1|$, then we get a $\mathbf{T}^2$ worth of valid limits for each domain. There is a non trivial requirement that needs to be satisfied for these limits to be glued together.

Apart from an easy use of an Arzela Ascoli type argument and elliptic bootstrapping, the main hurdle to proving Theorem 4.1 is dealing with the above types of non uniqueness for limits in showing that the different limiting pieces of our holomorphic curves glue together. One way to think of the problem at hand is that there will be a unique limit of our sequence of curves in the smooth part of $\mathbf{B}$, but there will be a family of valid tropical choices for our limit. We must construct a family of curves which contains all these tropical possibilities, and extend this family a little so that we can capture the behavior of our sequence of curves. There is some choice involved in this extension of a family, but we shall construct the family locally in coordinate charts. To make our choices match up, we shall use equivariant coordinate charts, which are constructed in Lemma A.3.

The following lemma constructs compatible $\mathbf{T}^n$ actions on neighborhoods of strata in $\mathbf{B}$. These $\mathbf{T}^n$ actions are defined on page 41. Locally, each $\mathbf{T}^n$ action is given by multiplying the $\mathbf{T}_P^n$ coordinates of $\mathbb{R}^m \times \mathbf{T}_P^n$ by constants, so

$$(\tilde{c}_1, \dots, \tilde{c}_n) * (x, \tilde{z}_1, \dots, \tilde{z}_n) = (x, \tilde{c}_1 \tilde{z}_1, \dots, \tilde{c}_n \tilde{z}_n)$$

We shall need these $\mathbf{T}^n$ actions to move parts of our curves around in a consistent way.

Lemma 4.3. *Given any basic exploded manifold $\hat{\mathbf{B}}$, and a family $\hat{\mathbf{B}} \longrightarrow \mathbf{G}$, for each strata $\hat{\mathbf{B}}_i \subset \hat{\mathbf{B}}$, there exists some open neighborhood $\mathbf{U}_i \subset \hat{\mathbf{B}}$ containing $\hat{\mathbf{B}}_i$ and a free action of $\mathbf{T}^n$ on $\mathbf{U}_i$ in the sense of definition A.2 (where n is the dimension of the tropical part of $\hat{\mathbf{B}}_i$). This $\mathbf{T}^n$ action on $\mathbf{U}_i$ satisfies the following properties:*

1. *The $\mathbf{T}^n$ action preserves fibers of the family $\hat{\mathbf{B}} \longrightarrow \mathbf{G}$, in the sense that if p_1 and p_2 have the same image in $\mathbf{G}$, then $\tilde{z} * p_1$ and $\tilde{z} * p_2$ also have the same image in $\mathbf{G}$.*
2. *If the closure of $\hat{\mathbf{B}}_i$ contains $\hat{\mathbf{B}}_j$, then the restriction of the action of $\mathbf{T}^n$ on $\mathbf{U}_i$ to $\mathbf{U}_i \cap \mathbf{U}_j$ is equal to the restriction of the $\mathbf{T}^m$ action on $\mathbf{U}_j$ to $\mathbf{U}_i \cap \mathbf{U}_j$ and to some subgroup.*

Proof:

This follows immediately from Lemma A.3. Choose equivariant coordinates on $\hat{\mathbf{B}} \rightarrow \mathbf{G}$, and let $\mathbf{U}_i$ be some open neighborhood of $\mathbf{B}_i$ contained inside the union of all our coordinate charts that intersect $\mathbf{B}_i$. As each of these coordinate charts intersect $\mathbf{B}_i$, and $\mathbf{B}$ is basic, there is a canonical inclusion of $\mathbf{B}_i$ into the tropical part of each of these coordinate charts, and a corresponding action of $\mathbf{T}^n$ on these coordinate charts given by multiplying the corresponding coordinates by constants. The fact that our coordinates are equivariant implies that this gives a $\mathbf{T}^n$ action on $\mathbf{U}_i$. The compatibility of the actions on $\mathbf{U}_i \cap \mathbf{U}_j$ and the compatibility of the action with the projection map $\hat{\mathbf{B}} \rightarrow \mathbf{G}$ all follow from the fact that these coordinate charts are equivariant. $\square$


Note the assumption that $\mathbf{B}$ is basic is mainly for convenience in the following arguments. In the family case, by restricting to a subset of $\mathbf{G}$ which contains the image of a subsequence, we can also assume that $\hat{\mathbf{B}}$ is basic.

We shall now start working towards proving Theorem 4.1. The proof shall rely on Lemma 3.14 on page 16 and Proposition 3.18 on page 21. We shall be using the notation from Proposition 3.18. First, choose the following:

1. Choose a metric, a taming form $\omega \in \Omega$, and finite collection of coordinate charts on $\mathbf{B}$ each contained in some $\mathbf{U}_i$ from Lemma 4.3 and small enough to apply Lemma 3.14. For the case of a family, restrict $\hat{\mathbf{B}} \rightarrow \mathbf{G}$ to some small coordinate chart on $\mathbf{G}$ in which the image of some subsequence converges, and choose our finite collection of coordinate charts on (the smaller, renamed,) $\hat{\mathbf{B}}$ satisfying the above.
2. Choose exploded annuli $\mathbf{A}_n^i \subset \mathbf{C}^i$ satisfying the conditions in Proposition 3.18 with ω -energy bound ϵ small enough and R large enough that each smaller annulus $\mathbf{A}_{\frac{6R}{10},n}^i$ is contained well inside some $\mathbf{U}_j$ in the sense that it is of distance greater than 2 to the boundary of $\mathbf{U}_j$. (The fact that we can achieve this follows from Lemma 3.12 on page 13.) Also choose ϵ small enough and R large enough so that each connected smooth component of $\mathbf{A}_{\frac{6R}{10},n}^i$ is contained in one of the above coordinate charts and can have Lemma 3.14 applied to it.

Use the notation C_m^i to indicate connected components of $\mathbf{C}^i - \bigcup_k \mathbf{A}_{\frac{8R}{10},k}^i$. We can choose a subsequence so that

- the number of such components is the same for each $\mathbf{C}^i$,
- C_m^i has topology that is independent of i ,
- and choosing diffeomorphisms identifying them, C_m^i converges as $i \rightarrow \infty$ to some C_m in the sense that the metric from Proposition 3.18 and the complex structure converges to one on C_m .



We shall view each of our annuli $\mathbf{A}_n^i$ as glued out of two semi infinite annuli, so $\mathbf{A}_n^i$ has two holomorphic coordinates $\tilde{z}^+$ and $\tilde{z}^-$ related by

$$\tilde{z}^+ \tilde{z}^- = Q_n^i \in \mathbb{R}^* \mathfrak{t}^{\mathbb{R}^+}$$

where the conformal modulus of $\mathbf{A}_n^i$ is $-\log Q_n^i$, and we regard $\mathbf{A}_n^i \subset (\mathbf{T}_1^1)^2$ as given by

$$\mathbf{A}_n^i = \{\tilde{z}^+ \tilde{z}^- = Q_n^i, |\tilde{z}^+| < 1, |\tilde{z}^-| < 1\} \subset (\mathbf{T}_1^1)^2$$

These Q_n will later become coordinates on our family $\mathbf{F}$.) Choose a subsequence so that the number of annuli is independent of i , and $\{Q_n^i\}$ converges. (In other words, calling any conformal modulus not a real number infinity, either the conformal modulus of $\mathbf{A}_n^i$ converges to a finite number, or converges to infinity.)

We can define


$$\mathbf{A}_n^\pm := \{|\tilde{z}^\pm| \leq e^{-\frac{7R}{10}}\} \subset \mathbf{T}_1^1$$

$$\mathbf{A}_{\frac{7R}{10},n}^i := \{(\tilde{z}^+, \tilde{z}^-) \text{ so that } \tilde{z}^+ \tilde{z}^- = Q_n^i\} \subset \mathbf{A}_n^+ \times \mathbf{A}_n^-$$

We shall need to keep track of what $\mathbf{A}_n$ is attached to, so use the notation $n^\pm$ to define $C_{n^\pm}^i$ as the component attached to the end of $\mathbf{A}_n^i$ with the boundary $|\tilde{z}_n^\pm| = 1$. (Assume, by passing to a subsequence, that $n^\pm$ is well defined independent of i .)

We can consider transition maps between $C_{n^\pm}^i$ and $\mathbf{A}_{\frac{7R}{10},n}^i$ to give transition maps between $C_{n^\pm}$ and $\mathbf{A}_n^\pm$. The bound on the derivative of transition maps from Proposition 3.18 on the region $\mathbf{A}_{\frac{9R}{10},n}^i - \mathbf{A}_{\frac{6R}{10},n}^i$ (and standard elliptic bootstrapping to get bounds on higher derivatives on the smaller region $\mathbf{A}_{\frac{8R}{10},n}^i - \mathbf{A}_{\frac{7R}{10},n}^i$) tells us that we can choose a subsequence so that the transition maps between $C_{n^\pm}^i$ and $\mathbf{A}_n^\pm$ converge to some smooth transition map between $C_{n^\pm}$ and $\mathbf{A}_n^\pm$. We can modify our diffeomorphisms identifying C_m^i with C_m so

that these transition maps all give exactly the same map. Denote the complex structure on C_m induced by this identification by j_i . (We shall do something similar for the annuli A_n^i later on.) We can do this so that j_i and the metrics given by this identification still converge in C^∞ to those on C_m .



We shall now start to construct our family $\hat{\mathbf{C}} \longrightarrow \mathbf{F}$. The first step shall be to get some kind of convergence for each piece of our holomorphic curve.

For each C_m choose some point $q_m \in C_m$, and consider

$$Q_m^i := f^i(q_m) \in \hat{\mathbf{B}}$$

This sequence Q_m^i has a subsequence that converges in $\hat{\mathbf{B}}$ to some point $Q_m \in \hat{\mathbf{B}}$. Label the strata of $\hat{\mathbf{B}}$ that contains Q_m by $\hat{\mathbf{B}}_m$, and consider the chart $\mathbf{U}_m$ from Lemma 4.3 containing $\hat{\mathbf{B}}_m$. As the derivative of f^i restricted to C_m is bounded, we can choose a subsequence so that $f^i(C_m)$ is contained well inside $\mathbf{U}_m$. (Note that any point in the strata $\hat{\mathbf{B}}_m$ is infinite distance to the boundary of $\mathbf{U}_m$. We can therefore choose a subsequence so that the image of $f^i(C_m)$ is contained in the subset of $\mathbf{U}_m$ which consists of all points some arbitrary distance from the boundary of $\mathbf{U}_m$.)

As the derivative of f^i is uniformly bounded on $\mathbf{C}^i - \bigcup A_{\frac{9R}{10},k}^i$, we can use lemmas 3.7, 3.8, and 3.9 to get bounds on the higher derivatives of f^i restricted to C_m^i . Lemma 3.1 then tells us that if $f^i(q_m)$ converges to $Q_m \in \hat{\mathbf{B}}$, the geometry around $f^i(q_m)$ converges to that around Q_m , so we can choose a subsequence so that f^i restricted to C_m^i converges in some sense to a map

$$f_{m,Q_m} : C_m \longrightarrow \hat{\mathbf{B}} \text{ so that } f_{m,Q_m}(q_m) = Q_m \subset \mathbf{U}_m$$

More specifically, remembering that everything is contained inside $\mathbf{U}_m$, we can use our $\mathbf{T}^n$ action on $\mathbf{U}_m$ to say this more precisely. There exists some sequence $\tilde{C}_m^i \in \mathbf{T}^n$ so that

$$\tilde{C}_m^i * f^i : C_m \longrightarrow \mathbf{U}_m$$


converges in C^∞ to $f_{m,Q_m} : C_m \longrightarrow \mathbf{U}_m$.

Of course, there was a choice involved here. We could also have chosen a different limit Q'_m of $f^i(q_m)$, which would give a translate of f_{m,Q_m} by our $\mathbf{T}^n$ action. These choices will turn up as parameters on our family. They will need to be ‘compatible’ in the sense that these pieces will need to fit together.

We now consider the analogous convergence on annular regions. Because the ω -energy of f^i restricted to our annular regions $\mathbf{A}_n^i$ is small, we can apply Lemma 3.14 on page 16 to tell us that if the limit of the conformal modulus of $\mathbf{A}_n^i$ is infinite, then f^i restricted to the smooth parts of $\mathbf{A}_n^i$ converges in some sense (considered in more detail later) to some unique pair of holomorphic maps

$$f_{n,Q}^\pm : \{e^{-\frac{7R}{10}} > |\tilde{z}_n^\pm| > 0\} \longrightarrow \hat{\mathbf{B}}$$

compatible with $f_{n^\pm, Q_{n^\pm}} : C_{n^\pm} \longrightarrow \hat{\mathbf{B}}$ and the transition maps. (Recall that $C_{n^\pm}$ is the component attached to $\mathbf{A}_n^\pm$.) As we have made the assumption that J is civilized, we can use the usual removable singularity theorem for finite energy holomorphic curves to see that these limit maps extend uniquely to smooth maps on $\{e^{-\frac{7R}{10}} > |\tilde{z}_n^\pm| > t^x\} \subset \mathbf{A}_n^\pm$ for some $x > 0$.



We may assume after passing to a subsequence that $f^i(\mathbf{A}_{\frac{6R}{10},n}^i)$ are all contained inside a single $\mathbf{U}_n$. Recalling our convention that $C_{n^\pm}$ intersects $\mathbf{A}_n^i$, note that this means that $\mathbf{U}_{n^\pm}$ intersects $\mathbf{U}_n$ (and $\mathbf{U}_{n^\pm} \subset \mathbf{U}_n$), so our sequence $\tilde{C}_{n^\pm}^i$ defined earlier also has an action on $\mathbf{U}_n$, and

$$\tilde{C}_{n^\pm}^i * f^i : \mathbf{A}_{\frac{7R}{10},n}^i \longrightarrow \mathbf{U}_n$$

converges in C^∞ on compact subsets adjacent to $C_{n^\pm}$ to $f_{n,Q}^\pm$.

We now have a type of convergence of the individual pieces we have cut our holomorphic curve into. We shall now define a model for the exploded structure on our family that is too large, as it ignores the requirement that these pieces must fit together. Later, we shall cut down this model to reflect the requirement of the pieces gluing together.

Define $\mathbf{V}_m \subset \mathbf{U}_m$ to be the exploded manifold consisting of all points $Q'_m \in \mathbf{U}_m$ so that

1. There exists some $\tilde{z}$ so that $Q'_m = \tilde{z} * Q_m$
2. Defining

$$f_{m,Q'_m} := \tilde{z} * f_{m,Q_m}$$

The image of C_m , $f_{m,Q'_m}(C_m)$ is contained well inside $\mathbf{U}_m$ in the sense that the distance to the boundary of $\mathbf{U}_m$ is greater than 1.

3. If C_m is attached to $\mathbf{A}_n^\pm$, then defining

$$f_{n,Q'}^\pm := \tilde{z} * f_{n,Q}^\pm$$

The image of the smooth part of $\mathbf{A}_n^\pm$, $\{f_{n,Q'}^\pm(\tilde{z}), 0 < |\tilde{z}| < e^{-\frac{7R}{10}}\}$ is also contained well inside $\mathbf{U}_n$.

Note that $\mathbf{V}_m$ is a smooth exploded manifold which includes all Q'_m which are limits for Q_m^i . For such Q'_m , the map f_{m,Q'_m} will be holomorphic, but for other Q'_m , this may not be the case.

We shall consider the family $\mathbf{F}$ as a sub exploded manifold of

$$\prod_m \mathbf{V}_m \prod_n \mathbf{Q}_n \subset \prod_m \mathbf{U}_m \prod_n \mathbf{Q}_n$$

The $\mathbf{Q}_n$ above stands for the ‘gluing parameter’ for identifying coordinates $\tilde{z}_n^\pm$ on $\mathbf{A}_n^\pm$ via

$$\tilde{z}_n^+ \tilde{z}_n^- = Q_n \in \mathbf{Q}_n := \{|\tilde{z}| < e^{-2R}\} \subset \mathbf{T}_1^1$$

We shall consider the following sequence of points

$$Q^i := (f^i(q_1), f^i(q_2), \dots, Q_{n_1}^i, \dots) \in \prod_m \mathbf{U}_m \prod_n \mathbf{Q}_n$$

where the conformal modulus of $\mathbf{A}_n^i$ is equal to $-\log Q_n^i$.

Note that there is a transitive action of $\mathbf{T}^k$ on $\prod \mathbf{V}_m \prod \mathbf{Q}_n$ which is the action from Lemma 4.3 on $\mathbf{V}_m \subset \mathbf{U}_m$, and multiplication by some coordinate of $\mathbf{T}^k$ on $\mathbf{Q}_n$. Our family will be given by a complete inclusion

$$\mathbf{F} \longrightarrow \prod_m \mathbf{V}_m \prod_n \mathbf{Q}_n \subset \prod_m \mathbf{U}_m \prod_n \mathbf{Q}_n$$

satisfying the following conditions:

1. The image of $\mathbf{F}$ contains some point $Q \in \prod_m \mathbf{V}_m \prod_n \mathbf{Q}_n$ which is a limit of Q^i , and the image of $\mathbf{F}$ is given by the orbit of Q under the action of some subgroup of $\mathbf{T}^k$.
2. The distance in any smooth metric from some subsequence of Q^i to the image of $\mathbf{F}$ converges to 0.
3. There is no other inclusion satisfying the above conditions which has smaller dimension than $\mathbf{F}$.

It is clear that such an $\mathbf{F}$ must exist. The entire space satisfies the first two properties, so we may simply choose an $\mathbf{F}$ satisfying the first two properties which has minimal dimension. It follows from the definition of an action of $\mathbf{T}^k$ on page 41 that $\mathbf{F}$ is smooth as it is the orbit of some $\mathbf{T}^n$ action. The minimality of the dimension of $\mathbf{F}$ is a trick which will ensure that our limiting maps $f^\pm$ defined on $\mathbf{A}^\pm$ glue together well. (See Lemma 4.4 below.)

Now let us construct our family of curves $\hat{\mathbf{C}} \longrightarrow \mathbf{F}$. We shall have charts on $\hat{\mathbf{C}}$ given by $\hat{C}_m := C_m \times \mathbf{F}$ and $\hat{\mathbf{A}}_n$. $\hat{\mathbf{A}}_n$ has coordinates $(\tilde{z}_n^+, \tilde{z}_n^-, Q)$ where $\tilde{z}_n^\pm \in \mathbf{A}_n^\pm$, $Q \in \mathbf{F}$ and $Q_n = \tilde{z}_n^+ \tilde{z}_n^-$. Transition maps between $\hat{C}_m$ and $\hat{\mathbf{A}}_n$ are simply given by the transition maps between C_m and $\mathbf{A}_n^\pm$ times the identity on the $\mathbf{F}$ component. For example, if ϕ is a transition map between C_m and $\mathbf{A}_n^+$, the corresponding transition map between $\hat{C}_m$ and $\hat{\mathbf{A}}_n$ is given by

$$(z, Q) \mapsto \left(\phi(z), \frac{Q_n}{\phi(z)}, Q \right)$$

This describes the exploded manifold $\hat{\mathbf{C}}$. The map down to $\mathbf{F}$ is simply given by the obvious projection to the last component of each of the above charts. This gives a smooth family of exploded curves.

Lemma 4.4. *If $Q \in \mathbf{F}$, and $Q_n^i \rightarrow Q_n$, then $f_{n,Q}^\pm$ can be glued by the identification $\tilde{z}_n^+ \tilde{z}_n^- = Q_n$. This is automatic if the limit of the conformal modulus of $\mathbf{A}_n^i$ is finite.*

Proof:

In the case that the limit of the conformal modulus of $\mathbf{A}_n^i$ is finite, $\lim Q_n^i = Q_n$ for any $Q \in \mathbf{F}$. If this was not true, we could simply restrict $\mathbf{F}$ to the points that satisfy this. This would include any point in $\mathbf{F}$ that is the limit of Q^i , it would also satisfy the other conditions required of $\mathbf{F}$, but have smaller dimension contradicting the minimality of $\mathbf{F}$. Therefore in the case that Q_n is finite as we have f^i converging on $C_{n^\pm}$ and $\mathbf{A}_n^i$, and transition maps between these also converging, we can glue the limit.

In the case that the limit of the conformal modulus of $\mathbf{A}_n^i$ is infinite, we want to glue $f_{n,Q}^+(\tilde{z}_n^+)$ to $f_{n,Q}^-(\tilde{z}_n^-)$ over the region where the smooth coordinates $[\tilde{z}_n^+] = [\tilde{z}_n^-] = 0$ via the identification $\tilde{z}_n^+ \tilde{z}_n^- = Q_n$. Define the following continuous function

$$\phi_n : \mathbf{F} \longrightarrow \mathbf{T}^l \text{ (the group acting on } \mathbf{U}_n \text{)}$$

which detects the failure for $f_n^\pm$ to glue for any $Q \in \mathbf{F}$: If $[Q_n] = 0$, and $\tilde{z}_n^+ \tilde{z}_n^- = Q_n$ so that $[\tilde{z}_n^+] = [\tilde{z}_n^-] = 0$, let $\phi(Q)$ be the element of $\mathbf{T}^l$ so that $\phi_n(Q) * f_{n,Q}^+(\tilde{z}_n^+) = f_{n,Q}^-(\tilde{z}_n^-)$. This may only fail to be defined because the left or right hand sides might not be inside $\mathbf{U}$. Note first that if defined, this doesn't depend on our choice of $\tilde{z}_n^\pm$. Also note that this will be defined on some open subset of $\mathbf{F}$, and there exists some homomorphism from $\mathbf{T}^k$ to $\mathbf{T}^l$ so that this map is equivariant with respect to the $\mathbf{T}^k$ action on $\mathbf{F}$ and the $\mathbf{T}^l$ action on $\mathbf{U}_m$. We can uniquely extend ϕ_n to the rest of $\mathbf{F}$ in an equivariant way.

We want to show that ϕ_n is identically 1 on $\mathbf{F}$. For each Q^i , there exists a close point $\check{Q}^i$ inside $\mathbf{F}$ so that the distance in any smooth metric between $\check{Q}^i$ and Q^i converges to 0 as $i \rightarrow \infty$. Then $\phi_n(\check{Q}^i)$ converges to 1, because these points Q^i come from holomorphic curves that are converging on either end of $\mathbf{A}_{\frac{7R}{10},n}^i$ to $f_n^\pm$ in a way given by Lemma 3.14. This tells us that the sub exploded manifold of $\mathbf{F}$ given by $\phi_n^{-1}(1)$ satisfies the conditions 1 and 2 above, so by the minimality of the dimension of $\mathbf{F}$, it must be all of $\mathbf{F}$. This tells us that if Q_n^{-1} is infinite, then we can glue together $f_n^\pm$ without any modifications. $\square$

Note that choosing any point $Q \in \mathbf{F}$ so that Q is a limit of Q^i (such a point must exist by the definition of $\mathbf{F}$), the above lemma allows us to glue together f_{m,Q_m} and $f_{n,Q}^\pm$ to obtain our limiting holomorphic curve f for Theorem 4.1. Note that condition 6 from Proposition 3.18 ensures that f is a stable holomorphic curve.

If the above lemma held for every point in $\mathbf{F}$, we would have constructed our family. As it is, we need to make some gluing choices.

If $Q_n \in \mathbb{C}^*$ and the limit of the conformal modulus of $\mathbf{A}_n^i$ is infinite, we will need to make 'gluing' choices. We do not use the standard gluing and cutting maps as this will not give us strong enough regularity for the resulting family.

1. Chose some sequence $\check{Q}^i$ of points in $\mathbf{F}$ so that the distance between $\check{Q}^i$ and Q^i converges to 0.

We need to take care of the different complex structures obtained by gluing $\mathbf{A}_n^\pm$ by $\check{Q}_n^i$ and Q_n^i . Choose an almost complex structure j_i on $\mathbf{A}_n^+$ as follows: Choose smooth isomorphisms

$$\Phi_n^i : \mathbf{A}_n^+ \longrightarrow \mathbf{A}_n^+$$

so that

(a)

$$\Phi_n^i(\tilde{z}_n^+) = \tilde{z}_n^+ \text{ for } |\tilde{z}_n^+| \geq e^{-\frac{8R}{10}}$$

(b)

$$\Phi_n^i(\tilde{z}_n^+) = \frac{\check{Q}_n^i}{Q_n^i} \tilde{z}_n^+ \text{ for } |\tilde{z}_n^+| \leq e^{-R}$$

- (c) On the region $[\tilde{z}_n^+] \neq 0$, the sequence of maps $\{\Phi_n^i\}$ converges in C^∞ to the identity.

Now define j_i on $\mathbf{A}_n^+$ to be the pullback under Φ_n^i of the standard complex structure. Using the standard complex structure on $\mathbf{A}_n^-$, we then get our j_i defined on $\hat{\mathbf{A}}_n$. This is compatible with the j_i already defined on $\hat{C}_m$, so we get j_i defined on $\hat{\mathbf{C}}$. Note that j_i restricted to the fiber over $\tilde{Q}^i$ is the complex structure on $\mathbf{C}^i$. From now on, we shall use these new coordinates on $\mathbf{A}_n^i$.

2. We now define the linear gluing map as follows:

- (a) Chose some smooth cutoff function

$$\rho : \mathbb{R}^* \longrightarrow [0, 1]$$

so that

$$\begin{aligned} \rho(x) &= 0 \text{ for all } x \geq e^{-\frac{8R}{10}} \\ \rho(x) &= 1 \text{ for all } x \leq e^{-\frac{9R}{10}} \end{aligned}$$

Extend this to

$$\rho : \mathbb{R}^* \mathfrak{t}^{\mathbb{R}} \longrightarrow [0, 1]$$

satisfying all the above conditions. (We defined this first on $\mathbb{R}^*$ so that it was clear what ‘smooth’ meant.)

- (b) Given maps $\phi^+, \phi^- : \mathbf{T}_1^1 \longrightarrow \mathbb{C}^k$ which vanish at $z = 0$ define the gluing map

$$G_{(\phi^+, \phi^-)}(\tilde{z}^+, \tilde{z}^-) := \rho(|\tilde{z}^-|) \phi^+(\tilde{z}^+) + \rho(|\tilde{z}^+|) \phi^-(\tilde{z}^-)$$

Note that if ϕ^+ and ϕ^- are smooth, $G_{(\phi^+, \phi^-)} : \mathbf{T}_2^2 \longrightarrow \mathbb{C}^k$ is smooth. Note also that if ϕ^+ and ϕ^- are small in $C^{\infty, \delta}$, then $G_{(\phi^+, \phi^-)}$ is small too.

3. We now define a linear ‘cutting’ map: as follows:

- (a) Choose a smooth cutoff function $\beta : \mathbb{R}^* \longrightarrow [0, 1]$ satisfying the following:

$$\begin{aligned} \beta(x) + \beta(x^{-1}) &= 1 \\ \beta(x) &= 1 \text{ for all } x > e^{\frac{R}{10}} \end{aligned}$$

- (b) Given a map $\phi : \mathbf{A}_n^i \longrightarrow \mathbb{C}^k$ and $\tilde{Q}_n^i \in \mathbb{C}^*$, where

$$\mathbf{A}_n^i := \{\tilde{z}^+ \tilde{z}^- = \tilde{Q}_n^i\} \subset (\mathbf{T}_1^1)^2$$

we can define the cutting of ϕ as follows:

$$\begin{aligned} \phi^+(\tilde{z}^+) &:= \beta\left(\frac{|z^+|^2}{|\tilde{Q}_n^i|}\right) \phi(z^+) \\ \phi^-(\tilde{z}^-) &:= \beta\left(\frac{|z^-|^2}{|\tilde{Q}_n^i|}\right) \phi(z^-) := \beta\left(\frac{|z^-|^2}{|\tilde{Q}_n^i|}\right) \phi\left(z^+ = \frac{\tilde{Q}_n^i}{z^-}\right) \end{aligned}$$

As $\phi^\pm(\tilde{z}^\pm)$ vanishes for $[\tilde{z}^\pm]$ small, we may extend $\phi^\pm$ to be zero for $\tilde{z}^\pm$ small.

Note that $G_{(\phi^+, \phi^-)}$ restricted to $\tilde{z}^+ \tilde{z}^- = \tilde{Q}_n^i$ is equal to ϕ .

4. Lemma 3.6 tells us that the image of our annuli of finite conformal moduli, $f^i(\mathbf{A}_{\frac{7R}{10},n}^i)$ is contained in some coordinate chart appropriate for Lemma 3.14. (We can choose this coordinate chart to be contained inside $\mathbf{U}_n$ so that the $\mathbf{T}^k$ action on $\mathbf{U}_n$ just consists of multiplying coordinate functions by a constant). In fact, we can choose a subsequence so that either $\mathbf{A}_n^i$ has infinite conformal modulus for all i or $f^i(\mathbf{A}_{\frac{7R}{10},n}^i)$ is contained in one of these coordinate charts for all i . In that case, Lemma 3.14 together with the conditions on our coordinate change for $\mathbf{A}_n^i$ above tells us that

$$\begin{aligned} f_n^i(z) &= (e^{\phi_{n,1}^i(z)} c_{n,1}^i z^{\alpha_1}, \dots, e^{\phi_{n,k}^i(z)} c_{n,k}^i z^{\alpha_k}, \dots, c_{n,d-k} + \phi_{n,d-k}^i) \\ &:= e^{\phi_n^i} c_n^i z^\alpha \end{aligned}$$

where ϕ_n^i is exponentially small on the interior of $\mathbf{A}_{R,n}^i$ as required by Lemma 3.14. If $Q^\infty \in \mathbf{F}$ is some limit of Q^i , then $f_{n,Q^\infty}^\pm$ in these coordinates are given by

$$\begin{aligned} f_{n,Q^\infty}^+(z^+) &:= e^{\phi_n^+(z^+)} c_n(z^+)^\alpha \\ &:= (e^{\phi_{n,1}^+(z^+)} c_{n,1}(z^+)^{\alpha_1}, \dots, e^{\phi_{n,k}^+(z^+)} c_{n,k}(z^+)^{\alpha_k}, \dots, c_{n,d-k} + \phi_{n,d-k}^+(z^+)) \\ f_{n,Q^\infty}^-(z^-) &:= e^{\phi_n^-(z^-)} c_n \left(\frac{Q_n^\infty}{z^-} \right)^\alpha \end{aligned}$$

where $\phi^\pm$ are smooth and vanish when $[z^\pm] = 0$.

Similarly, if $\mathbf{A}_n^i$ is infinite, then we can consider f_n^i to map to $\mathbf{O}_n$ so that

$$\begin{aligned} f_n^i(z^+) &:= e^{\phi_n^{i+}(z^+)} c_n^i(z^+)^\alpha \\ f_n^i(z^-) &:= e^{\phi_n^{i-}(z^-)} c_n^i \left(\frac{\check{Q}_n^i}{z^-} \right)^\alpha \end{aligned}$$

5. If $p_1 = c * p_2$ using one of our $\mathbf{T}^n$ actions, use the notation $c = \frac{p_1}{p_2}$, so $p_1 = \frac{p_1}{p_2} * p_2$. Define $F_n^i : \hat{\mathbf{A}}_n \longrightarrow \hat{\mathbf{B}}$ by

$$F_n^i(\tilde{z}^+, \tilde{z}^-, Q) := \frac{Q_{n^+}}{\check{Q}_{n^+}^i} * c_n^i(\tilde{z}^+)^\alpha$$

and

$$F_n(\tilde{z}^+, \tilde{z}^-, Q) := \frac{Q_{n^+}}{Q_{n^+}^\infty} * c_n(\tilde{z}^+)^\alpha$$

(Recall that Q_{n^+} recorded the position of C_{n^+} which is attached to $\mathbf{A}_n^+$).

Then we can define $\hat{f}^i : \hat{\mathbf{A}}_n \longrightarrow \hat{\mathbf{B}}$ using the cutting and gluing maps above by

$$\hat{f}^i(\tilde{z}^+, \tilde{z}^-, Q) := e^{G_{(\phi_n^{i+}, \phi_n^{i-})}(\tilde{z}^+, \tilde{z}^-)} F_n^i(\tilde{z}^+, \tilde{z}^-, Q)$$

We can similarly define $\hat{f} : \hat{\mathbf{A}}_n \longrightarrow \hat{\mathbf{B}}$.

Also define

$$\hat{f} : \hat{C}_m \longrightarrow \hat{\mathbf{B}}$$

by

$$\hat{f}(z, Q) = f_{m, Q_m}(z)$$

and define $\hat{f}^i$ on $\hat{C}_m$ similarly by translating f^i appropriately depending on the difference between $\tilde{Q}_m^i$ and Q_m , so

$$\hat{f}^i(z, Q) = \frac{Q_m}{\tilde{Q}_m^i} * f_m^i(z)$$

This gives well defined smooth maps

$$\hat{f}^i : \hat{\mathbf{C}} \longrightarrow \hat{\mathbf{B}}$$

$$\hat{f} : \hat{\mathbf{C}} \longrightarrow \hat{\mathbf{B}}$$

so that f^i is given by the restriction of $\hat{f}^i$ to the fiber of $\hat{\mathbf{C}} \longrightarrow \mathbf{F}$ over $\tilde{Q}^i \in \mathbf{F}$. Note that as the vectorfields $\phi^\pm$ are vertical with respect to $\hat{\mathbf{B}} \longrightarrow \mathbf{G}$, G_{ϕ^+, ϕ^-} is too, and we have finally constructed our required families.

$$\begin{array}{ccc} (\hat{\mathbf{C}}, j_i) & \xrightarrow{\hat{f}^i} & \hat{\mathbf{B}} \\ \downarrow & & \downarrow \\ \mathbf{F} & \longrightarrow & \mathbf{G} \end{array} \quad \begin{array}{ccc} (\hat{\mathbf{C}}, j) & \xrightarrow{\hat{f}} & \hat{\mathbf{B}} \\ \downarrow & & \downarrow \\ \mathbf{F} & \longrightarrow & \mathbf{G} \end{array}$$

Lemma 4.5.

$$\hat{f}^i : (\hat{\mathbf{C}}, j_i) \longrightarrow \hat{\mathbf{B}}$$

converges in $C^{\infty, 1}$ to

$$\hat{f} : (\hat{\mathbf{C}}, j) \longrightarrow \hat{\mathbf{B}}$$

Proof:

We work in coordinates. First note that $\hat{f}^i$ converges in C^∞ restricted to $\hat{f}$ on $\hat{C}_m$. This also holds for $\hat{\mathbf{A}}_n$ if the size of $\mathbf{A}_n^i$ stays bounded. Note also that j_i converges in C^∞ to j .

It remains only to show that $\hat{f}^i$ converges in $C^{\infty, \delta}$ to $\hat{f}$ on $\hat{\mathbf{A}}_n$. Recall that $\hat{\mathbf{A}}_n$ has coordinates $(\tilde{z}^+, \tilde{z}^-, Q)$ where $Q \in \mathbf{F}$ and $Q_n = \tilde{z}^+ \tilde{z}^-$. Note that the maps $F_n^i : \hat{\mathbf{A}}_n \longrightarrow \hat{\mathbf{B}}$ defined above converge in C^∞ to $F_n : \hat{\mathbf{A}}_n \longrightarrow \hat{\mathbf{B}}$. We have that

$$\hat{f}^i(\tilde{z}^+, \tilde{z}^-, Q) := e^{G_{(\phi_n^{i+}, \phi_n^{i-})}(\tilde{z}^+, \tilde{z}^-)} F_n^i(\tilde{z}^+, \tilde{z}^-, Q)$$

Similarly, we write

$$\hat{f}(\tilde{z}^+, \tilde{z}^-, Q) := e^{G_{(\phi_n^+, \phi_n^-)}(\tilde{z}^+, \tilde{z}^-)} F_n(\tilde{z}^+, \tilde{z}^-, Q)$$

We can choose a subsequence so that $\phi_n^{i\pm}(z^\pm)$ converges in C^∞ on compact subsets of $\{0 < |z^\pm| \leq e^{-\frac{7R}{10}}\}$ to $\phi_n^\pm(z^\pm)$. Lemma 3.14 and our cutting construction above tells us that for any $\delta < 1$, there exists some constant c independent of n so that

$$|\phi_n^{i\pm}(\tilde{z}^\pm)| \leq c |\tilde{z}^\pm|^\delta$$

and the same inequality holds for $\phi_n^\pm$ and all derivatives (with a different constant c). This implies that $\phi_n^{i\pm}$ converges to $\phi_n^\pm$ in $C^{\infty,1}$. Adding extra coordinates on which a function does not depend does not affect $C^{\infty,1}$ convergence. Therefore $\phi_n^{i\pm}(\tilde{z}^\pm)$ converges in $C^{\infty,1}$ to $\phi_n^\pm(\tilde{z}^\pm)$ on $\hat{\mathbf{A}}_n$. Then $G_{(\phi_n^{i+}, \phi_n^{i-})}$ is obtained by adding these two functions multiplied by fixed smooth function cutoff functions. Therefore, $G_{(\phi_n^{i+}, \phi_n^{i-})}$ converges in $C^{\infty,1}$ to $G_{(\phi_n^+, \phi_n^-)}$ on $\hat{\mathbf{A}}_n$. This implies that $\hat{f}^i$ converges to $\hat{f}$ in $C^{\infty,1}$ as required. $\square$

This completes the proof of Theorem 4.1.

A Equivariant coordinates

In this appendix, we prove that any exploded manifold has coordinate charts with transition functions in a special form.

Suppose that $\mathbb{R}^n \times \mathbf{T}_P^m$ and $\mathbb{R}^{n'} \times \mathbf{T}_Q^{m'}$ are two coordinate charts on $\mathbf{B}$ which intersect non trivially and that P is a face of the polytope Q . By making a monomial change of coordinates, we may also assume that the coordinates on P are the first m coordinates on Q , and that the remaining $m' - m$ coordinates on Q vanish on P . Then the transition map between $\mathbb{R}^n \times \mathbf{T}_P^m$ and $\mathbb{R}^{n'} \times \mathbf{T}_Q^{m'}$ must be in the form

$$\phi(\tilde{x}, \tilde{z}) = (f, g_1 \tilde{z}_1, \dots, g_m \tilde{z}_m, g_{m+1}, \dots, g_{m'})$$

where

$$\begin{aligned} f : \mathbb{R}^n \times [\mathbf{T}_P^m] &\longrightarrow \mathbb{R}^{n'} \\ g_i : \mathbb{R}^n \times [\mathbf{T}_P^m] &\longrightarrow \mathbb{C}^* \end{aligned}$$

are smooth functions. A special case is when f and g_i depend only on $\mathbb{R}^n$. We shall prove that any exploded manifold $\mathbf{B}$ can be covered with coordinate charts modeled on open subsets of $\mathbb{R}^n \times \mathbf{T}_P^m$ with transition maps in this special form.

Definition A.1. Define a $(\mathbb{C}^*)^m$ action on $\mathbb{R}^n \times \mathbf{T}_P^m$ by

$$(c_1, \dots, c_m) * (x, \tilde{z}) = (x, c_1 \tilde{z}_1, \dots, c_m \tilde{z}_m)$$

Similarly, if $(\tilde{c}_1, \dots, \tilde{c}_m) \in \mathbf{T}^m$,

$$\text{let } (\tilde{c}_1, \dots, \tilde{c}_m) * (x, \tilde{z}) = (x, \tilde{c}_1 \tilde{z}_1, \dots, \tilde{c}_m \tilde{z}_m) \text{ where defined}$$

Call a map f from a subset of $\mathbb{R}^n \times \mathbf{T}_P^m$ to $\mathbb{R}^{n'} \times \mathbf{T}_Q^{m'}$ equivariant if there exists a map

$$\alpha : \mathbf{T}^m \longrightarrow \mathbf{T}^{m'}$$

so that

$$f(c * (x, \tilde{z})) = \alpha(c) * f(x, \tilde{z})$$

whenever the left hand side is defined.

Note that every map from $\mathbb{R}^n \times \mathbf{T}_P^m$ is equivariant when restricted to the interior of P .

Definition A.2. A free action of $\mathbf{T}^k$ on an exploded manifold $\mathbf{B}$ is a map from a subset $\text{Dom}(\ast)$ of $\mathbf{T}^k \times \mathbf{B}$ to $\mathbf{B}$

$$\begin{aligned} \text{Dom}(\ast) &\longrightarrow \mathbf{B} \\ (\tilde{z}, p) &\mapsto \tilde{z} \ast p \end{aligned}$$

so that

1. There exists a covering of $\mathbf{B}$ by coordinates modeled on open subsets of $\mathbb{R}^n \times \mathbf{T}_P^m$ so that the $\mathbf{T}^k$ action is given by multiplication on the last k coordinates of $\mathbf{T}_P^m$, in the sense that
 - (a) if p and $\tilde{z} \ast p$ are in the same coordinate chart, then $\tilde{z} \ast p$ is determined by multiplying the last k coordinates of p by $\tilde{z}$.
 - (b) If q is a point in the coordinate chart obtained from p by multiplying the last k coordinates by $\tilde{z}$, then $(\tilde{z}, p) \in \text{Dom}(\ast)$ and $\tilde{z} \ast p = q$.
2. If $(\tilde{z}, p) \in \text{Dom}(\ast)$, then $(\tilde{w}, \tilde{z} \ast p) \in \text{Dom}(\ast)$ if and only if $(\tilde{w}\tilde{z}, p) \in \text{Dom}(\ast)$, and in this case,

$$\tilde{w} \ast (\tilde{z} \ast p) = (\tilde{w}\tilde{z}) \ast p$$

Note that an action of $\mathbf{T}^n$ on $\mathbf{B}$ is not quite an action of $\mathbf{T}^n$ considered as a group, simply because the action may not be defined everywhere.

Lemma A.3. Every exploded manifold $\mathbf{B}$ has a cover by coordinate charts modeled on open subsets of $\mathbb{R}^n \times \mathbf{T}_P^m$ so that every transition map is either equivariant or its inverse is equivariant.

Every family $\pi : \mathbf{B} \longrightarrow \mathbf{G}$ of exploded manifolds has a cover by coordinate charts so that transition maps on $\mathbf{G}$ and $\mathbf{B}$ are equivariant in the above sense, and π is equivariant in the sense that given any of our coordinate charts U on $\mathbf{B}$, one of our coordinate charts V on $\mathbf{G}$ contains $\pi(U)$, and $\pi : U \longrightarrow V$ is equivariant.

Proof:

Suppose that we have already constructed some coordinate charts with equivariant transition maps which cover an open subset $O \subset \mathbf{B}$.

Let $\psi : V \longrightarrow \mathbb{R}^n \times \mathbf{T}_{P^\circ}^m$ be a coordinate chart which restricts to a coordinate chart $\psi^\circ : U \longrightarrow \mathbb{R}^n \times \mathbf{T}_{P^\circ}^m$ on a strata $\mathbf{B}_i$ of $\mathbf{B}$, where $P^\circ \subset P$ is the interior of P , and U is $V \cap B_i$. This coordinate chart ψ° is automatically compatible with all previous coordinate charts because P° is an open polytope. We shall now try to extend ψ° to a coordinate chart $\psi' : V' \longrightarrow \mathbb{R}^n \times \mathbf{T}_P^m$ on an open neighborhood V' of U in $\mathbf{B}$. We shall not necessarily be able to make the transition maps between this new chart and our previous charts equivariant, but if we choose any open subset O' of $\mathbf{B}$ so that the closure of O' is contained in O , then our new coordinate chart can have equivariant transition maps to the old coordinate charts restricted to O' .

Denote by O_U the set of points $p \in O \cap V$ so that there exists some previously defined coordinate chart containing p so that $\tilde{z} \ast p \in U$. Clearly, O_U is open and contains $U \cap O$. Therefore,

$$V' := (V \setminus \bar{O}') \cup O_U$$

is an open subset of V which contains U .

On O_U there is an action of $\mathbf{T}^n$. We can define another map $\psi' : O_U \longrightarrow \mathbb{R}^n \times \mathbf{T}_P^m$ so that $\psi' = \psi$ restricted to U and $\psi'(\tilde{z} * p) = \tilde{z} * \psi'(p)$. The new map ψ' is an isomorphism onto its image because ψ was. The goal is now to extend ψ' to V' . There exists a smooth vectorfield v on $\psi(O_U)$ so that v vanishes on U , and ψ' is given by ψ composed with exponentiating v using the standard flat metric on $\mathbb{R}^n \times \mathbf{T}_P^m$. To extend ψ' , we shall extend this vectorfield. Of course, not every vectorfield defined on an open subset extends, but we can extend v restricted to $O' \cap O_U$ by multiplying v by a cutoff function which is 1 on O' and 0 on $V' \setminus O_U$. The resulting map given by composing ψ with exponentiation by v will be a local isomorphism on a neighborhood of U in V' , so by restricting to this smaller neighborhood if necessary, we get our required coordinate chart ψ' which is compatible with all previous coordinate charts restricted to O' .

This proof shall now follow by a double induction. Suppose that every strata of $\mathbf{B}$ with dimension strictly less than k in $[\mathbf{B}]$ can be covered by coordinate charts with equivariant transition maps (this is trivially true for $k=0$). We shall prove that we can cover all strata of dimension less or equal than k in $[\mathbf{B}]$ by coordinate charts with equivariant transition maps. Assume that we already have our equivariant cover of the strata of lower dimension. Let O denote the open set already covered, and let O' be an open subset of O which contains all strata of lower dimension so that the closure of O' is contained in O . Then choose a sequence of open sets O_i containing O' so that $O_0 = O$ and the closure of O_{i+1} is contained inside O_i .

Denote by U_0 the intersection of O' with the strata of dimension k in $[\mathbf{B}]$. Choose a countable open cover $\{U_i\}$ of the strata of dimension k in $[\mathbf{B}]$ so that for all $i > 0$, $\bar{U}_i$ is contained inside some coordinate chart. Assume for our second induction that for all $i < n$, there exists a coordinate chart V_i containing the closure of some open neighborhood V'_i of U_i so that all transition maps between V_i for $i \leq n$ and all previous coordinate charts are equivariant restricted to some open neighborhood N of $O_{n-1} \bigcup_{i=1}^{n-1} \bar{V}'_i$.

Choose some open neighborhood N' of $O_n \bigcup_{i=1}^{n-1} \bar{V}'_i$ whose closure is contained in N . The first section of this proof showed that we can construct a coordinate chart V_n which contains the closure of some open neighborhood V'_n of U_n so that all transition maps between V_n and all previous coordinate charts are equivariant restricted to N' . Therefore, our second inductive statement holds for all n . Now consider our previously constructed coordinate charts restricted to O' together with the new coordinate charts V'_i . The transition maps between these coordinate charts are all equivariant, and together these coordinate charts cover all strata of dimension k in $\mathbf{B}$. By induction, we may cover all of $\mathbf{B}$ by coordinate charts with equivariant transition maps.

We can prove the family case similarly. First choose equivariant coordinate charts on $\mathbf{G}$. We can modify any coordinate chart on $\mathbf{B}$ so that π is equivariant. The difference between π on a given coordinate chart and π on an equivariant coordinate chart is given by the flow of a time dependent vector field on $\mathbf{G}$, we can lift this vector field to a time dependent vector field on our coordinate chart and use the flow of this vectorfield to modify our coordinate chart so that π is equivariant. We may then repeat the above argument for $\mathbf{B}$ choosing all coordinate charts so that the map π is equivariant. The step where a coordinate chart needs to be modified using the flow of a vector field now uses only vertical

vectorfields, so this procedure preserves the property that π is equivariant. $\square$

B A symplectic analogue of the explosion functor

The following example constructs a symplectic exploded manifold from a symplectic manifold with ‘normal crossing symplectic divisors’. The construction is analogous to the explosion functor, but is not functorial.

Example B.1.

Suppose that we have a compact symplectic manifold M with some collection of codimension 2 embedded symplectic submanifolds N_i which intersect each other orthogonally. (In other words, if $x \in N_i \cap N_j$, the vectors $v \in T_x M$ so that $\omega(v, \cdot)$ vanishes on $T_x N_i$ are contained inside $T_x N_j$.)

We shall construct an exploded manifold with an almost complex structure tamed by a symplectic form from the above data as follows

1. Let I index a set of submanifolds, and let

$$N_I := \bigcap_{i \in I} N_i$$

2. Suppose that for all $|I| \geq k$ we have achieved the following:

- (a) A neighborhood of N_I is identified with the sum of $|I|$ complex line bundles over N_I with complex linear connections ∇ .

$$\begin{array}{c} \mathbb{C}^{|I|} \longrightarrow E_I \\ \downarrow \\ N_I \end{array}$$

- (b) The symplectic structure restricted to the zero section of E_I is standard in the $\mathbb{C}^k$ directions, and the $\mathbb{C}^k$ directions are symplectically orthogonal to N_I .
- (c) If $i \in I$, then the intersection of N_i with our neighborhood of N_I corresponds to the subset of E_I where the i th line bundle coordinate is 0.
- (d) If $I' \supset I$, then on some neighborhood of $N_{I'}$, the bundle $E_I \longrightarrow N_I$ is equal to the $\mathbb{C}^I$ bundle with total space $E_{I'}$ and projection map the map that sends the coordinates corresponding to $I \subset I'$ to 0.

$$\begin{array}{c} \mathbb{C}^{|I|} \longrightarrow E_{I'} \\ \downarrow \\ N_I \cap E_{I'} \end{array}$$

This identification preserves the connections ∇

- (e) There is an almost complex structure J tamed by ω on a neighborhood of all N_I with $|I| \geq k$ so that J restricted to the fibers of E_I is the standard complex structure, N_I is a J holomorphic submanifold, and J on ∇ -horizontal subspaces of TE_I is the lift of J from TN_I .
3. Now assuming the inductive hypothesis 2, we shall show that the same can be achieved for $|I| = k - 1$.

Over a neighborhood of $\bigcap_{I' \supset I} N_{I'} \subset N_I$, the compatibility condition 2d ensures that we have $|I|$ complex line bundles locally given by the total space $E_{I'}$ and the projection map which sends the coordinates corresponding to $I \subset I'$ to 0.

$$\begin{array}{c} \mathbb{C}^{|I|} \longrightarrow E_{I'} \\ \downarrow \\ N_I \cap E_{I'} \end{array}$$

These complex line bundles satisfy the requirement 2b and come with connections and an almost complex structure satisfying the requirement 2e. There is no obstruction to choose an identification of a neighborhood of N_I with the total space E_I of $|I|$ line bundles satisfying the conditions 2b, 2c and 2e. The fact that 2b and 2c can be satisfied simultaneously follows from the requirement that our symplectic submanifolds N_i intersect symplectically orthogonally. To satisfy requirement 2e, choose an almost complex structure J on N_I tamed by ω which agrees with the previously defined J on some neighborhood of $\bigcap_{I' \supset I} N_{I'} \subset N_I$, then extend the definition of J to a neighborhood of N_I by giving the fibers of E_I the standard complex structure and lifting J to horizontal subspaces of TE_I from TN_I . The fact that such a J agrees on some neighborhood of N_I with the previously defined J follows from the compatibility condition 2d. This almost complex structure J is tamed by ω restricted to some neighborhood of N_I because of the condition 2b which requires that at the zero section the symplectic structure on the fibers is standard, and that the fibers are symplectically orthogonal to N_I .

4. Therefore, as the inductive hypothesis 2 holds for k larger than the number of submanifolds, we get that we can achieve 2 for all k . Now cover M with coordinate charts $\mathbb{R}^n \times \mathbb{C}^k$, so that such a coordinate is contained inside E_I for some $I = k$ and the $\mathbb{C}^k$ coordinate gives coordinates for the k complex line bundles in E_I . Replace these coordinates (x, z) with coordinates $(x, \tilde{z}) \in \mathbb{R}^n \times (\mathbf{T}_1^1)^n$. We can replace transition maps similarly to the explosion functor. In particular, if the old transition map was ϕ , then the new transition map $\tilde{\phi}$ can be defined as follows:

$$\begin{aligned} x \circ \tilde{\phi} &= x \circ \phi \circ [\cdot] \\ \text{if } z_i \circ \phi &= f(x)z_j \text{ then } \tilde{z}_i \circ \tilde{\phi} = f(x)\tilde{z}_j \\ \text{if } z_i \circ \phi &\text{ is nonvanishing, then } \tilde{z}_i \circ \tilde{\phi} = z_i \circ \phi \circ [\cdot] \end{aligned}$$


These new coordinates and transition maps define a basic exploded manifold $\mathbf{M}$ with smooth part $[\mathbf{M}] = M$.

5. There is a smooth almost complex structure J on $\mathbf{M}$ compatible with the smooth part map and the almost complex structure on M . We can define this almost complex structure in coordinates as follows: In the old coordinate charts, J exchanged the real and imaginary parts of $z \frac{\partial}{\partial z}$, therefore, in our new coordinate charts, we can get J to exchange the real and imaginary parts of $\tilde{z} \frac{\partial}{\partial \tilde{z}}$. In our old coordinate charts,

$$\left(J \frac{\partial}{\partial x_i} \right) z_j = f(x) z_j$$

where $f : \mathbb{R}^n \rightarrow \mathbb{C}$ is smooth. This is because J was defined using the lift of an almost complex structure on $\mathbb{R}^n$ using k complex linear connections. Therefore, $\left(J \frac{\partial}{\partial x_i} \right)$ is a sum of smooth functions times $\frac{\partial}{\partial x_i}$ and the real and imaginary parts of $z \frac{\partial}{\partial z}$. In our new coordinate charts, we can define $\left(J \frac{\partial}{\partial x_i} \right)$ to be the corresponding sum of smooth functions times $\frac{\partial}{\partial x_i}$ and the real and imaginary parts of $\tilde{z} \frac{\partial}{\partial \tilde{z}}$. The pullback of ω to $\mathbf{M}$ via the smooth part map $\mathbf{M} \rightarrow M$ is a symplectic form which tames the almost complex structure J .

The following is a picture of the result of this procedure on a toric symplectic manifold, where for our symplectic submanifolds we use the fixed loci of circle actions. We shall represent the smooth part $[\mathbf{M}]$ by its moment map:



It is interesting to notice that in this toric case, the dual fan to the moment map of M is the tropical part $\underline{\mathbf{M}}$.

C Construction of a strict taming

Lemma C.1. *Suppose that an exploded manifold $\mathbf{B}$ satisfies the following:*

1. $\mathbf{B}$ is basic and complete.
2. All strata of the tropical part $\underline{\mathbf{B}}$ have the integral affine structure of either a standard simplex or $\mathfrak{t}^{[0,\infty)^n}$

Then given any civilized almost complex structure tamed by a symplectic form ω on $\mathbf{B}$, there exists a close by civilized almost complex structure J with a strict taming Ω which contains ω .

This new almost complex structure can be chosen C^0 close to the old almost complex structure considered as an almost complex structure on $[\mathbf{B}]$, and $C^{\infty,1}$ close when considered as an almost complex structure on $\mathbf{B}$. In the special case that $\mathbf{B}$ is a complex exploded manifold, no modification of the complex structure is required. Another case in which no modification is required is when for any exploded function $\tilde{z}$ defined on a subset of $\mathbf{B}$, $\tilde{z}^{-1}\bar{\partial}\tilde{z}$ is a one form which is equal to a smooth one form pulled back from $[\mathbf{B}]$.

Proof:

Lemma A.3 implies that we may choose coordinate charts with equivariant transition maps in the sense that after some monomial change of coordinates, transition maps or their inverses are in the form

$$(x, \tilde{z}_1, \dots, \tilde{z}_m) \mapsto (f(x), g_1(x)\tilde{z}_1, \dots, g_m(x)\tilde{z}_m, g_{m+1}(x), \dots, g_{m'}(x))$$

We shall label strata of $\mathbf{B}$ by the polytope which is the closure of their image in the tropical part of $\mathbf{B}$. Each strata $\mathbf{B}_P$ is in the form of some $\mathbf{T}_{P^\circ}^m$ bundle over $[\mathbf{B}_P]$, where P° is the interior of P and $[\mathbf{B}_P]$ is the interior of a smooth compact symplectic manifold (M_P, ω) with a collection of positively intersecting codimension 2 embedded symplectic submanifolds M_Q . (There is one such M_Q for each polytope Q in $\underline{\mathbf{B}}$ which has P as a codimension 1 face). By interior we mean the complement of these symplectic submanifolds. Using the above coordinates with equivariant transition maps, we may identify a neighborhood $\mathbf{U}_P$ of $\mathbf{B}_P$ with an open subset of a $\mathbf{T}_P^m$ bundle over $[\mathbf{B}_P]$.

Starting with the polytopes P of maximal dimension, construct functions

$$r_{F \subset P} : \mathbf{U}_P \longrightarrow \mathbb{R}^* \mathfrak{t}^{[0,\infty)}$$

for each codimension 1 face F of P satisfying the following conditions.

1. In our coordinates, $(x, \tilde{z})$ on $\mathbb{R}^n \times \mathbf{T}_P^m$, if F is given by $\tilde{z}_k = \mathfrak{t}^0$, then

$$r_{F \subset P}(x, \tilde{z}) = f(x) |\tilde{z}_k|$$

and if F is given by $\tilde{z}_1, \dots, \tilde{z}_m = \mathfrak{t}^c$, then

$$r_{F \subset P}(x, \tilde{z}) = f(x) \mathfrak{t}^c |\tilde{z}_1^{-1} \dots \tilde{z}_m^{-1}|$$

(In the above, f is a smooth function which may depend on F .)


2. If P is a standard simplex of size c , then

$$\prod_F r_{F \subset P} = 1 \mathfrak{t}^c$$

3. If Q is a face of P , and $F \cap Q$ is a codimension 1 face of P , then on $\mathbf{U}_Q \cap \mathbf{U}_P$,

$$r_{F \subset P} = r_{(F \cap Q) \subset Q}$$

The diagram below may help with visualizing $r_{F \subset P}$. The picture of $[\mathbf{T}_P^2]$ could be understood as a moment map of a $\mathbb{T}^2$ action, in which case $r_{F \subset P}$ will actually be the square root of distance in the directions indicated below.



On each of our coordinate charts on $\mathbf{U}_P$, there is an action of $\mathbf{T}^m$ given by multiplying the coordinates of $\mathbf{T}_P^m$ by the corresponding constants in $\mathbf{T}^m$. Our transition maps are equivariant with respect to this action, so there is a $\mathbf{T}^m$ action on $\mathbf{U}_P$ in the sense of definition A.2. We may consider $\mathbf{U}_P$ as an open subset of a corresponding $\mathbf{T}_P^m$ bundle with the same $\mathbf{T}^m$ action, and $r_{F \subset P}$ defined similarly. If our original almost complex structure is not integrable, we may modify our almost complex structure to an almost complex structure J which is preserved by the $\mathbf{T}^m$ action. By restricting our coordinate charts to be smaller if necessary, we can make this new almost complex structure $C^{\infty,1}$ close to the old almost complex structure on $\mathbf{B}$, and C^0 close to the old almost complex structure on $[\mathbf{B}]$. Therefore, this modification of our complex structure can be chosen small enough that ω still tames J . As $\mathbf{B}$ is complete, we may assume (by adding extra coordinates obtained by restricting those on adjacent strata if necessary) that there exists some $r_0 > 0$ so that $\mathbf{U}_P$ contains the subset of this $\mathbf{T}_P^m$ bundle where all $r_{F \subset P}$ are less than r_0 .

Define a one form $\alpha_{F \subset P}$ on $\mathbf{U}_P$ for each codimension one face F of P as follows:

$$\alpha_{F \subset P} := r_{F \subset P}^{-1} dr_{F \subset P} \circ J$$

In the case that we modified our almost complex structure, the one form $\alpha_{F \subset P}$ is preserved by the action of $\mathbf{T}^m$ on $\mathbf{U}_P$, so $d\alpha_{F \subset P}$ can be regarded as the pullback of a form on $[\mathbf{B}_p]$. In the case that J is complex, we may consider $\alpha_{F \subset P}$ in holomorphic coordinate charts. After a monomial change of coordinates, $r_{F \subset P} = f |\tilde{z}_1|$, where f is some smooth nonvanishing real valued function on our holomorphic coordinate chart. Then $d\alpha_{F \subset P} = d(f^{-1} df \circ J)$, which is a two form which is pulled back from the smooth part of the coordinate chart.

If modification of J is undesirable, it is not necessary so long as any $d\alpha_{F \subset P}$ is the pullback of a two form on $[\mathbf{B}]$, which is equivalent to the requirement that for any exploded function $\tilde{z}$ locally defined on $\mathbf{B}$, the two form $d(\tilde{z}^{-1} d\tilde{z} \circ J)$ is a smooth two form pulled back from $[\mathbf{B}]$. It is sufficient that $\tilde{z}^{-1} \bar{\partial} \tilde{z}$ is a smooth one form pulled back from $[\mathbf{B}]$. It is easy to check that our modified J and any integrable complex structure satisfies this condition. The set of J tamed by ω satisfying this condition is convex.

Note that if Q is a face of P and $F \cap Q$ is a codimension 1 face of Q , the fact that $r_{F \subset P} = r_{(F \cap Q) \subset Q}$ implies that


$$\alpha_{F \subset P} = \alpha_{(F \cap Q) \subset Q} \text{ on } \mathbf{U}_P \cap \mathbf{U}_Q$$

We now provide a strict taming Ω which for each point $p \in \mathbf{B}$ includes some two form ω_p taming J well in a neighborhood of p as in part 2 of the above definition of a strict taming. We do this as follows: Suppose that $p \in \mathbf{U}_P$ and $r_{F \subset P}(p)$ is much less than r_0 for all codimension 1 faces of P and $r_{P \subset Q}$ is not too small for all strata Q with P as a codimension 1 face. We will specify the location of p more precisely later.

Consider the set $S_p \subset \mathbb{R}^* \mathfrak{t}^{\mathbb{R}}$ given by

$$S_p := \{r_{F \subset P}(p)\}$$

We shall define ω_p on a refinement of $\mathbf{B}$ determined by a subdivision of the tropical part $\underline{\mathbf{B}}$. Subdivide the polytope P by the set of all planes of the form $\{r_{F \subset P} \in \underline{S}_p\}$. (This is the set of planes through p parallel to the boundary of the strata and their image under any symmetry of the strata.) Similarly, if Q is connected to P through polytopes in $\underline{\mathbf{B}}$ with nonzero dimension, then subdivide Q by the planes $\{r_{F \subset Q} \in \underline{S}_p\}$. This subdivision of $\underline{\mathbf{B}}$ defines a refinement of $\mathbf{B}$.



P

We can now define a two form ω_p on the refinement of $\mathbf{B}$ defined above as follows: First choose a function

$$\phi : \mathbb{R}^* \mathfrak{t}^{\mathbb{R}} \longrightarrow \mathbb{R}$$

so that

$$\phi(r) = 0 \quad \forall r > 2\mathfrak{t}^0$$

$$\phi(r) + \phi(r^{-1}) = -1$$

and considered as a function $\phi : \mathbb{R}^* \longrightarrow \mathbb{R}$, ϕ is smooth, increasing and


$$\frac{d\phi(r)}{dr} > \frac{1}{r} \quad \text{for all } \frac{1}{\sqrt{2}} < r < \sqrt{2}$$

Also choose some smooth function $\rho : \mathbb{R}^* \mathfrak{t}^{[0, \infty)} \longrightarrow \mathbb{R}$ so that $\rho(r) = 0$ for all $r \geq r_0$ and $\rho(r) = 1$ for all $r < \frac{r_0}{2}$.

Now define $h : \mathbb{R}^* \mathfrak{t}^{\mathbb{R}^+} \longrightarrow \mathbb{R}$ by

$$h(r) := c\rho(r) + \frac{c}{|S_p|} \sum_{x \in S_p} \phi(rx^{-1})$$

where c is a positive constant that we shall choose later.



On $\mathbf{U}_P$, replace ω with

$$\omega_p := \omega + \sum_F d(h(r_{F \subset P})\alpha_{F \subset P}))$$

Now we do the same for all strata $\mathbf{B}_Q$ connected to $\mathbf{B}_P$ through strata with tropical part of non zero dimension. So on $\mathbf{U}_Q$,

$$\omega_p := \omega + \sum_F d(h(r_{F \subset P})\alpha_{F \subset P}))$$

For all other $\mathbf{U}'_P$, we simply leave ω unchanged. The fact that we chose the $r_{F \subset P}$ and $\alpha_{F \subset P}$ compatibly implies that the above construction defines ω_p as a smooth two form on our refinement. Actually, there exists a smooth two form on a refinement of $\mathbf{B} \times \mathbf{U}_P$ so that ω_p is the restriction of this two form to $\mathbf{B} \times p$. As ω_p differs from ω by d of a one form, the integral of ω agrees with the integral of ω_p over any complete exploded curve.

We must now consider whether ω_p is non negative on holomorphic planes, and whether it satisfies condition 2. On $\mathbf{U}_Q$ can break up ω_p into

$$\omega_p = \omega_0 + \omega_1$$

where

$$\begin{aligned} \omega_0 &:= \omega + c \sum_F d(\rho(r_{F \subset Q})\alpha_{F \subset Q}) + \frac{c}{|S_p|} \sum_{x \in S_p} \sum_F \phi(x^{-1}r_{F \subset Q})d\alpha_{F \subset Q} \\ \omega_1 &:= \frac{c}{|S_p|} \sum_{x \in S_p} \sum_F d(\phi(x^{-1}r_{F \subset Q})) \wedge \alpha_{F \subset Q} \end{aligned}$$

where c is the positive constant in the definition of h which we have yet to specify. Let us choose $c > 0$ to be a constant so that

1. On M_Q (the manifold with interior $[\mathbf{B}_Q]$), the two form

$$\omega + c \sum d(\rho(r_{F \subset Q'})\alpha_{F \subset Q'})$$


is still positive on holomorphic planes, where the sum involves all terms that are defined. Note that on $\mathbf{U}_{Q'}$, $d\alpha_{F \subset Q'}$ can be considered as a two form pulled back from the smooth part of the coordinate chart. It follows that $d(\rho(r_{F \subset Q'})\alpha_{F \subset Q'})$ is a two form on M_Q wherever $r_{F \subset Q'}$ is defined, even though $\alpha_{F \subset Q'}$ can't be considered as a one form on M_Q .

2. On M_Q , the two form $\omega + \sum_F x_i d\alpha_{F \subset Q}$ is still positive on holomorphic planes for all choices of x_i between 0 and c .

If we make these choices, then ω_0 will be nonnegative on any holomorphic plane, and positive on any holomorphic plane containing a vector with nonzero image in $\mathbf{P}$. As for ω_1 , on the complex subspace which is the intersection of $\ker dr_{F \subset Q}$ and $J(\ker dr_{F \subset Q})$ for all F , this two form ω_1 vanishes. As ϕ is monotone increasing, ω_1 is also nonnegative on the complimentary complex subspace which in coordinates is spanned by the real and imaginary parts of $\tilde{z}_i \frac{\partial}{\partial \tilde{z}_i}$. Therefore, ω_1 is also nonnegative on holomorphic planes, so ω_p is nonnegative

on holomorphic planes. Moreover, in a neighborhood of p in our refinement of $\mathbf{B}$, ω_p is positive on holomorphic planes. Therefore, restricting to any compact subset of some neighborhood of $\mathbf{B}_P$, we can obtain a uniform estimate in the form of 2. Then as we can cover all of $\mathbf{B}$ with compact subsets of neighborhoods of $\mathbf{B}_i$, we get a uniform estimate of the required form 2. $\square$

The following is a picture of the above procedure carried out in the case that $\mathbf{B}$ is the explosion of a toric manifold. We shall draw the smooth part of $\mathbf{B}$ using the moment map from ω and $\mathbf{B}'$ using the moment map from ω_p .



References

- [1] Frederic Bourgeois, Yakov Eliashberg, Helmut Hofer, Kris Wysocki, and Eduard Zehnder. Compactness results in symplectic field theory. *Geom. Topol.*, 7:799–888 (electronic), 2003.
- [2] M. Gromov. Pseudoholomorphic curves in symplectic manifolds. *Invent. Math.*, 82(2):307–347, 1985.
- [3] Eleny-Nicoleta Ionel and Thomas H. Parker. The symplectic sum formula for Gromov-Witten invariants. *Ann. of Math. (2)*, 159(3):935–1025, 2004.
- [4] An-Min Li and Yongbin Ruan. Symplectic surgery and Gromov-Witten invariants of Calabi-Yau 3-folds. *Invent. Math.*, 145:151–218, 2001.
- [5] Jun Li. A degeneration formula of GW-invariants. *J. Differential Geom.*, 60(2):199–293, 2002.
- [6] Dusa McDuff and Dietmar Salamon. *J-holomorphic Curves and Symplectic Topology*, volume 52 of *Colloquium Publications*. American Mathematical Society, 2004.
- [7] Brett Parker. Exploded manifolds. arXiv:0910.4201, 2009.
- [8] Brett Parker. Holomorphic curves in exploded torus fibrations: Regularity. arXiv:0902.0087v1, 2009.